

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
1 March 2001 (01.03.2001)

PCT

(10) International Publication Number
WO 01/14420 A2(51) International Patent Classification⁷: C07K 14/00

the University of California, 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). TAMAGNONE, Luca [IT/IT]; Corso Einaudi, 43, I-10129 Torino (IT).

(21) International Application Number: PCT/US00/23365

(22) International Filing Date: 25 August 2000 (25.08.2000)

(74) Agent: COX, Niki, D.; Biogen, Inc., 14 Cambridge Center, Cambridge, MA 02142 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/150,576 25 August 1999 (25.08.1999) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(71) Applicants (*for all designated States except US*): UNIVERSITY OF TORINO [IT/IT]; Department of Biomedical Sciences and Human Oncology, IRCC, SP 142, I-10060 Candiolo (IT). REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US).(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ARTIGIANI, Stefania [IT/IT]; Corso Brunelleschi, 121/B, I-10100 Torino (IT). COMOGLIO, Paolo, M. [IT/IT]; Strada Valsalice, 183/8, I-10100 Torino (IT). GOODMAN, Corey, S. [US/US]; Regents of the University of California, 12th floor, 1111 Franklin Street, Oakland, CA 94607-5200 (US). TESIER-LAVIGNE, Marc [US/US]; Regents of

Published:

*Without international search report and to be republished upon receipt of that report.**For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: NOVEL PLEXINS AND USES THEREOF

(57) Abstract: The invention provides methods and compositions related to novel plexins. The polypeptides may be produced recombinantly from transformed host cells and from the disclosed plexin encoding nucleic acids or purified from human cells. The invention provides isolated plexin hybridization probes and primers capable of specifically hybridizing with the disclosed plexin genes, plexin-specific binding agents such as specific antibodies, and methods of making and using the subject compositions in diagnosis, therapy and in biopharmaceutical industry. The invention also provides novel plexin neuropilin multimeric receptor complexes for semaphorins and methods of use thereof, including but not limited to, the treatment and diagnosis of neurological disease and neuroregeneration, immune modulation, and viral and oncological diseases.



WO 01/14420 A2

NOVEL PLEXINS AND USES THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

- 5 The invention relates to the identification and characterization of four novel proteins that are members of the plexin family.

Summary of the Related Art

- Plexin family members encode large transmembrane proteins, whose cysteine-rich extracellular domains share regions of homology with the Scatter Factor receptors (encoded by the Met gene family). The extracellular domains of plexins also contain ~500 amino acid Semaphorin domains (see below). The highly conserved cytoplasmic moieties of plexins (approx. 600 amino acids), however, have no homology with the Met tyrosine kinase domain, nor with any other known protein. Met-like receptors and their ligands, the Scatter Factors, mediate a complex biological program including
- 15 dissociation of cell-cell contacts, motility and invasion (for a review see Tamagnone, L. and Comoglio, P.M. (1997) "Control of invasive growth by hepatocyte growth factor (HGF) and related scatter factors." Cytokine Growth Factor Rev 8, 129-142). During embryogenesis Scatter Factor-1 and Met promote the dissociation of cell layers in the somites and drive the migration of myogenic cells to their appropriate location (Bladt, F., Riethmacher, D., Isenmann, S., Aguzzi, A., and Birchmeier, C. (1995) "Essential
- 20 role for the c-met receptor in the migration of myogenic precursor cells into the limb bud." Nature 376, 768-771; Maina, F., Casagrande, F., Audero, E., Simeone, A., Comoglio, P.M., Klein, R.a., and Ponzetto, C. (1996) "Uncoupling of grb2 from the met receptor in vivo reveals complex roles in muscle development." Cell 87, 531-542).
- 25 Met and Scatter Factor-1 are also involved in controlling neurite outgrowth and axonal guidance (Ebens, A., Brose, K., Leonardo, E.D., Hanson, M.G.J., Bladt, F., Birchmeier, C., Barres, B.A., and Tessier-Lavigne, M. (1996) "Hepatocyte growth factor/scatter factor is an axonal chemoattractant and a neurotrophic factor for spinal motor neurons." Neuron 17, 1157-1172; Maina, F., Hilton, M.C., Ponzetto, C., Davies, A.M., and
- 30 Klein, R. (1997) "Met receptor signaling is required for sensory nerve development and HGF promotes axonal growth and survival of sensory neurons." Genes Dev 11, 3341-3350; Maina, F., Hilton, M.C., Andres, R., Wyatt, S., Klein, R., and Davies, A.M.

(1998) "Multiple roles for hepatocyte growth factor in sympathetic neuron development." *Neuron* 20, 835-846).

The first clue regarding a possible function for plexins came from the finding that a novel plexin, *Vespr*, interacts with the viral semaphorin A39R (Comeau, M.R., Johnson, R., DuBose, R.F., Petersen, M., Gearing, P., VandenBos, T., Park, L., Farrah, T., Buller, R.M., Cohen, J.I., Strockbine, L.D., Rauch, C., and Spriggs, M.K. (1998) "A poxvirus-encoded semaphorin induces cytokine production from monocytes and binds to a novel cellular semaphorin receptor, *VESPR*." *Immunity* 8, 473-482). Semaphorins are a large family of secreted and membrane-bound molecules that are characterized by an extracellular domain containing a ~500 amino acid Semaphorin domain (Kolodkin et al. (1993) "The semaphorin genes encode a family of transmembrane and secreted growth cone guidance molecules." *Cell* 75, 1389-1399). As noted above, plexins contain a more divergent but nevertheless conserved Semaphorin domain.

Semaphorins were originally characterized in the nervous system, where they have been implicated in repulsive axon guidance (Kolodkin et al. (1993) *supra*; Luo, Y., Raible, D., and Raper, J.A. (1993) "Collapsin: a protein in brain that induces the collapse and paralysis of neuronal growth cones." *Cell* 75, 217-227; Tessier-Lavigne, M. and Goodman, C.S. (1996) "The molecular biology of axon guidance." *Science* 274, 1123-1133). More recently, semaphorins have been furthermore implicated in cardiac and skeletal development (Behar, O., Golden, J.A., Mashimo, H., Schoen, F.J., and Fishman, M.C. (1996) "Semaphorin III is needed for normal patterning and growth of nerves, bones and heart." *Nature* 383, 525-528), in the immune response (Hall, K.T., Boumsell, L., Schultze, J.L., Boussiotis, V.A., Dorfman, D.M., Cardoso, A.A., Bensussan, A., Nadler, L.M., and Freeman, G.J. (1996) "Human CD100, a novel leukocyte semaphorin that promotes B-cell aggregation and differentiation." *Proc.Natl.Acad.Sci.U.S.A.* 93, 11780-11785), in the regulation of angiogenesis (Miao, H.Q., Soker, S., Feiner, L., Alonso, J.L., Raper, J.A., and Klagsbrun, M. (1999) "Neuropilin-1 mediates collapsin-1/Semaphorin III inhibition of endothelial cell motility. Functional competition of collapsin-1 and vascular endothelial growth factor-165" [In Process Citation]. *J Cell Biol* 146, 233-242), and in tumor growth and metastasis (Christensen, C.R., Klingelhofer, J., Tarabykina, S., Hulgaard, E.F., Kramerov, D., and Lukanidin, E. (1998) "Transcription of a novel mouse semaphorin

gene, M-semaH, correlates with the metastatic ability of mouse tumor cell lines." Cancer Res. 58, 1238-1244).

Previously identified plexins have been shown to be expressed in the developing nervous system, (i.e. Plexin-A is a receptor for class 1 semaphorins (Sema-1a and Sema-1b). Moreover, Plexin-A has been shown via genetic analysis to control motor and CNS axon guidance induced by semaphorins (Winberg, M.L., Noordermeer, J.N., Tamagnone, L., Comoglio, P.M., Spriggs, M.K., Tessier-Lavigne, M., and Goodman, C.S. (1998). Plexin A is a neuronal semaphorin receptor that controls axon guidance. Cell 95, 903-916).

Thus a need exists for discovery of other members of the plexin family of proteins.

SUMMARY OF THE INVENTION

The present invention provides four novel plexin family members.

In one aspect, the invention provides an isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in (SEQ ID NO: 2), (SEQ ID NO: 4), (SEQ ID NO: 6) and (SEQ ID NO: 8).

In other aspects, the invention provides isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown in (SEQ ID NO: 1), (SEQ ID NO: 3), (SEQ ID NO: 5) and (SEQ ID NO: 7).

In another aspect, the invention provides a vector comprising the nucleic acid of the above-aspects.

The invention also provides an isolated polypeptide the amino acid sequence of which comprises residues 1-18, 19-518, 451-530, 601-680, 751-830, 800-1010, or 1196-1215 of SEQ ID NO: 2; 1-23, 24-507 or 1100-1119 of SEQ ID NO: 4; or 1-42, 43-600, 541-620, 691-770, 831-910, 900-1110 or 1270-1293 of SEQ ID NO: 6; or 8-49, 154-199 or 1-199 of SEQ ID NO: 8.

In another aspect, the invention provides an isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in (SEQ ID NO: 2), (SEQ ID NO: 4), (SEQ ID NO: 6) and (SEQ ID NO: 8)

The invention also provides a chimeric molecule comprising a polypeptide of the above aspects fused to a heterologous amino acid sequence. In one embodiment the heterologous amino acid sequence is a Fc region of an immunoglobulin.

In other aspects, the invention provides an antibody that specifically
5 binds to the polypeptides of the above aspects.

The invention also provides a method of treating, suppressing or altering a disorder involving aberrant immune regulation involving a signaling pathway between a plexin and a neuropilin in a mammal comprising the step of administering an effective amount of an agent to said mammal capable of interfering with the association between
10 the plexin and neuropilin. Contemplated agents include a chimeric molecule or an antibody of the above aspects.

DESCRIPTION OF THE DRAWINGS

Figure 1.

(A) Phylogenetic tree of human plexins. Known family members cluster in two
15 major groups: plexin A and plexin-B subfamilies. (B) Structural features of plexins, Met-like receptors and semaphorins. In the extracellular moieties, yellow boxes indicate the "sema" domains and blue boxes mark the cysteine-rich MRS motifs, some of which are stippled to indicate their atypical sequence; atypical MRS are also found in the *sema* domain of semaphorins. Sequence identity among *sema* domains ranges from 15-50%,
20 as previously described (see Winberg et al., 1998 *supra*). Potential furin-like proteolytic sites are marked by red ribbons. plexin-B1 "truncated" is the product of a splicing variant (see text). plexin-D1 and plexin-C1 (VESPR) are more distant family members, since they include atypical features in their extracellular domains. The intracellular domain of plexins (SP domain) is highly conserved through all family
25 members, and it includes two separate regions of high homology (Maestrini, E., Tamagnone, T., Longati, P., Cremona, O., Gulisano, M., Bione, S., Tamanini, F., Neel, B.G., Toniolo, D., and Comoglio, P.M. (1996) "A family of transmembrane proteins with homology to the MET-hepatocyte growth factor receptor." Proc. Natl. Acad. Sci. USA 93, 674-678) (green oval and box). Met-like receptors are disulfide-bound
30 heterodimers and include a cytoplasmic tyrosine kinase domain (red box). Mammalian semaphorins can be either secreted or cell surface proteins. Molecular weights of representative proteins are Plexin-A1 220 kDa, Plexin-B1 250 kDa, Plexin-C1 200

kDa, HGF-R/Met 145+45 kDa (heterodimer), Sema 4D 150 kDa (transmembrane), Sema7A approx. 100 kDa (membrane GPI-linked).

Figure 2.

(a) Cell surface semaphorins specifically bind human plexins. Micrographs of the binding assays done testing i) the extracellular domain of semaphorin CD100 fused to alkaline phosphatase (CD100-AP) on COS cells transfected with *plexin-B1 cDNA*; ii) control AP on plexin-B1; iii) CD100-AP on plexin-B2; iv) CD100-AP on the entire extracellular domain of plexin-B1; v) CD100-AP on isolated "plexin-B1 truncated" (including *sema* domain, 1° and 2° MRS); vi) CD100-AP on a "plexin-B1-Δsema" (including 2° and 3° MRS; vii) extracellular domain of semaphorin A39R fused to AP, on plexin-C1 (Vespr); viii) SemaK1-AP on plexin-C1. The final detection of the binding was done either using alkaline phosphatase substrates (i-vi) or by immunofluorescence (vii and viii). (B) Scatchard analysis and binding curve of CD100-AP to plexin-B1 ($K_D = 0.9 \text{ nM} \pm 0.15$).

Figure 3.

Plexins associate with neuropilins *via* specific extracellular domains. Western blots of immunoprecipitated samples from cells co-expressing neuropilins and plexins. Specific MoAbs were used, directed against the VSV-tag included in plexins or the myc-tag included in neuropilin-2 (Np2, 130 kDa). Np2 co-immunoprecipitates with plexins, such as plexin-A3 (220 kDa), the extracellular domain of plexinA1 (approx. 160 kDa), and plexin-B1 (250 kDa) but not with the unrelated cell surface receptor DCC (170 kDa). Np2 can also associate a truncated form of the extracellular moiety of plexin-B1 ("plex-B1 trunc.", approx. 110 kDa), containing the *sema* domain.

Figure 4.

Expression of mRNAs for plexins A1 (panel A, B), -A2 (panel C, D) and A3 (panel E, F) in the spinal cord (sc), dorsal root ganglia (d) and sympathetic ganglia (sg) of E13.5 mouse embryos. Expression of the mRNAs was detected by RNA in situ hybridization. Scale bar: 1 μm .

Figure 5.

Effect of a truncated plexin-A1 construct (lacking the cytoplasmic domain) on repulsive and attractive responses of *Xenopus* spinal neurons to Sema3A and netrin-1. (A-F) A control spinal neuron exposed to a gradient of Sema3A emanating from a pipette (A) is repelled away over a period of 1 hr (B). In contrast, a GFP-expressing

spinal neuron from an embryo injected with mRNA for the truncated plexin-A1 construct (C) is not affected by Sema3A (D). A similar neuron (E) shows a normal attractive response to netrin-1 (F).

(G) Cumulative distribution plot of turning angles for all the neurons studied.

- 5 Curves show the percent of neurons with turning angles less than the angle indicated on the abscissa, under different conditions (open circles, control neurons; black and blue circles, control neurons responding to Sema3A or netrin-1, respectively; red and green circles, responses of neurons expressing the truncated plexin-A1 construct to Sema3A and netrin-1, respectively. (H) Mean turning angle under all the conditions just
10 mentioned.

Figure 6.

- Tyrosine phosphorylation of plexin-A3 and plexin-B1. (a) Anti-phosphotyrosine western blotting of immunoprecipitated p220^{plex-A3} and p250^{plex-B1} proteins. plexin-B1 is larger since it contains an extra sequence between the second and
15 the third MRS motif, in the extracellular domain (see Fig. 1). (b) The same immunoprecipitated samples underwent *in vitro* kinase assay in the presence of [γ -³²P]ATP, Mg⁺⁺ and Mn⁺⁺ ions. The SDS-PAGE was treated with alkali in conditions known to eliminate the phosphate labeling of Ser/Thr residues and specifically preserving phosphotyrosines.

Figure 7

- Plexin-A3 overexpression mediates cell repelling cues. (a) Epithelial kidney MDCK cells transfected to overexpress plexin-A3 (or mock transfected) were cocultured with mesenchymal KJ-29 or NIH-3T3 cells. After 16-30 hours, mixed cultures of control cells (upper panels) reached confluency and stopped growing:
25 typically the epithelial cells formed islets (circled) surrounded by a fibroblasts lawn. In contrast, MDCKs overexpressing plexin-A3 (lower panels) overwhelmed the adjacent mesenchymal cells. The latter withdrew and selectively detached from the culture dish (dying cell clusters are indicated by arrowheads), and eventually epithelial cells only survived. To allow an easier detection of the mesenchymal cells, these were labeled
30 with DiI before being plated in mixed cultures. (b) Plexin-A3 expressing cells do not induce apoptotic signal on repelled fibroblasts. Mixed cultures of NIH 3T3 and control or plexin-A3 overexpressing MDCKs were tested for the presence of TUNEL-AP positive cells. Apoptotic cells were not present in clusters of repelled cells (indicated by

arrows). The right panel shows a positive control where apoptosis was induced on the same cells by UV treatment. (c) Plexin-A3 over-expressing cells form very transient contacts with fibroblasts. Time-lapse video-microscopy of control and plexin-A3 overexpressing MDCK cells grown in presence of fibroblasts. On top, snap-shot images from the movie, taken every 50 minutes (real time). In the upper row is shown the persistent contact of a fibroblast (marked by an arrow) with an islet of control MDCK cells (marked by a star). In the lower row another fibroblast, instead, forms a transient contact with an islet of plexin-A3 transfected cells, which also, in turn, reshapes. At the bottom, the diagrams show the relative frequency of persistent, transient or very transient contacts between fibroblasts and MDCK cells.

DETAILED DESCRIPTION OF THE INVENTION

The reference works, patents, patent applications, and scientific literature, including accession numbers to GenBank database sequences, that are referred to herein establish the knowledge of those with skill in the art and are hereby incorporated by reference in their entirety to the same extent as if each was specifically and individually indicated to be incorporated by reference. Any conflict between any reference cited herein and the specific teachings of this specification shall be resolved in favor of the later. Likewise, any conflict between an art-understood definition of a word or phrase and a definition of the word or phrase as specifically taught in this specification shall be resolved in favor of the latter.

Four novel human plexins have been identified: plexin-B2, plexin-B3, plexin-D1 and Plexin A-4. Plexin-A4 is located on human chromosome 7 and is a family member of the plexin-A subfamily which also includes plexin-A1 (alternatively named plexin-1/NOV), plexin-A2 (alternatively named plexin-2/OCT) and plexin-A3 (alternatively named plexin-2/SEX). Plexin-B2 and plexin-B3 are located on human chromosome 22 and chromosome X, respectively, and are family members of the plexin-B subfamily which also includes plexin-B1 (alternatively named SEP). Plexin-B3 maps very close to the plexin-A3 genomic locus on Xq28. Plexin-D1 is the first identified member of the plexin-D subfamily and is atypical of any of the other subfamilies. A fourth subfamily of plexins, the plexin-C subfamily, is defined by VESPR (now plexin-C1).

The four novel plexins as described herein have in their extracellular domains regions of homology with two other protein families: (a) Scatter Factors Receptors, encoded by the *MET* oncogene family (Tamagnone and Comoglio, 1997 *supra*), and (b)

Semaphorins (Kolodkin et al. (1993) *supra* (Figure 1b). In particular, plexins and Met-like receptors contain short cysteine-rich motifs, termed "Met Related Sequences" (MRS), whose minimal consensus is: C-X(5-6)-C-X(2)-C-X(6-8)-C-X(2)-C-X(3-5)-C (Maestrini et al., 1996 *supra*); Tamagnone and Comoglio, 1997 *supra*); blue boxes in Fig. 1B). The proteins of the Met family contain a single MRS (in their receptor β chains), whereas in plexin family members there are two/three repeated MRS motifs. Furthermore, all plexin family members contain in their extracellular moiety a 500 amino acid region similar to the sema domain of semaphorins (Kolodkin et al. (1993) *supra*; Winberg et al., 1998 *supra*); yellow boxes in Fig. 1B. The MRS motif is proposed to function as protein-protein interaction domain.

The cytoplasmic domain of plexins contains a ~600 amino acid domain which we term the SP domain ("Sex and Plexins", marked in green in Fig. 1B) that is highly conserved within the family (57-97% similarity) and in evolution (over 50% similarity between invertebrates and humans). The SP domain does not share homology with any known protein. It includes a number of potential tyrosine phosphorylation sites, but lacks the typical motifs of catalytic tyrosine kinases. Interestingly, the predicted secondary structure of the SP domain includes long conserved alpha helices, typically found in protein-protein interaction modules. Furthermore, there are several dihydrophobic amino acid motifs (such as LL or LI), known to mediate the internalization and downregulation of transmembrane receptors (Sandoval, I.V. and Bakke O. (1994). Targeting of membrane proteins to endosomes and lysosomes. Trends in Cell Biology 4, 292-297).

The present invention also demonstrates that plexins can form complexes with neuropilins, which in turn demonstrates that a receptor for semaphorins can be hetero-oligomers of plexins and neuropilins. As demonstrated by in situ mRNA expression analysis, plexins and neuropilins are in fact simultaneously expressed in several neuronal populations during embryonic development. The plexin-neuropilin complex predates ligand binding, since the association is not influenced by the presence of class 3 semaphorins. That the observed plexin-neuropilin complexes are formed in *cis* is furthermore supported by the experimental conditions used (cotransfection of isolated cells with the two constructs). An interaction in *trans* might also be envisioned (considering that plexins and semaphorins share similar *sema* domains), however by

analyzing mixed cultures of cells separately transfected with plexins and neuropilins we did not isolate associated complexes (data not shown).

We observed that the main semaphorin binding domain of neuropilins (CUB domain (Giger, R.J., Urquhart, E.R., Gillespie, S.K., Levengood, D.V., Ginty, D.D., and Kolodkin, A.L. (1998) "Neuropilin-2 is a receptor for semaphorin IV: insight into the structural basis of receptor function and specificity." *Neuron* 21, 1079-1092; Nakamura, F., Tanaka, M., Takahashi, T., Kalb, R.G., and Strittmatter, S.M. (1998) "Neuropilin-1 extracellular domains mediate semaphorin D/III-induced growth cone collapse" [In Process Citation]. *Neuron* 21, 1093-1100; Chen et al. 1998 *supra*) is not required for the interaction with plexins, as indicated by the association of the relevant Neuropilin-2 deletion construct with plexin-B1 (not shown). A ternary complex, where neuropilins use two distinct protein modules to form a bridge between the sema domain of semaphorins and the sema domain of plexins is thus contemplated. It is further contemplated that plexins are the functional partners of neuropilins, required for transducing signals mediated by semaphorins, preferably class 3 semaphorins. For example, in flies, which lack both neuropilins and class 3 semaphorins, D Plex A appears sufficient as a functional receptor for Sema 1a, a transmembrane class 1 semaphorin (Winberg et al., 1998 *supra*). Further support that plexins are functional co-receptors for secreted semaphorins is demonstrated in an experiment that shows that a truncated plexin-A1 construct expressed in *Xenopus* spinal neurons abolishes repulsive responses to Sema3A without markedly affecting attractive responses to netrin-1. These results are consistent with the involvement of plexins.

The intracellular signals transduced by plexins are still largely obscure. The cytoplasmic domain of plexins is large and highly conserved within and across species and contains stretches of alpha helices, which are putative protein-protein interaction domains, and could thus mediate the association with cytosolic partners. We demonstrate herein that the cytoplasmic domain of plexins can be tyrosine phosphorylated, suggesting that, like other receptors devoid of intrinsic catalytic activity, plexins may signal by associating a tyrosine kinase (Stahl, N. and Yancopoulos, G.D. (1993). The alphas, betas, and kinases of cytokine receptor complexes. *Cell* 74, 587-590; Glass, D.J., Bowen, D.C., Stitt, T.N., Radziejewski, C., Bruno, J., Ryan, T.E., Gies, D.R., Shah, S., Mattsson, K., Burden, S.J., DiStefano, P.S.,

Valenzuela, D.M., DeChiara, T.M., and Yancopoulos, G.D. (1996). Agrin acts via a MuSK receptor complex. *Cell* 85, 513-523).

In addition, we show herein that expression of plexins, particularly plexin-A3, mediates cell-repelling cues. By time-lapse video-microscopy we observed a true
5 repelling effect on fibroblasts. Intriguingly, we observed that -upon interaction with fibroblasts- also the islets of plexin-A3 MDCKs at times reshaped. This may be explained by the existence of intra-epithelial repelling cues, balanced by the attractive forces exerted by epithelial cell junctions.

Moreover we have demonstrated that in the nervous system (i.e. *Drosophila*), that
10 defasciculating motor axons co-express both Plexin A and one of its interacting partners, the transmembrane semaphorin Sema-1a (Winberg et al., 1998 *supra*). This demonstrates that plexins act *in vivo* either as receptors or ligands for cell surface semaphorins, which in turn can transduce intracellular signals, as reported for ephrins (Holland et al., 1996 *supra*). Semaphorins, therefore, besides being pivotal in axon
15 guidance, have a general role in morphogenesis and tissue remodeling by mediating cell-repelling cues via their interactions with plexins.

Accordingly, in a first aspect, the invention provides an isolated nucleic acid molecule encoding a novel human plexin polypeptide. By "plexin polypeptide" is meant a member of the plexin family comprising an amino acid sequence that shares at least
20 60% amino acid sequence homology with SEQ ID NOS: 2 (plexin B-2), 4 (plexin B-3), 6 (plexin D-1) or 8 (plexin A-4), preferably, at least 65% sequence homology, more preferably, at least 70% sequence homology, more preferably, at least 75% sequence homology, more preferably, at least 80% sequence homology, still more preferably at least 85% sequence homology, even more preferably, at least 90% sequence
25 homology, and most preferably at least 95% sequence homology with SEQ ID NOS: 2, 4, 6 or 8. Plexin polypeptides of the invention are useful for modulating cell growth (i.e. nerve) and immune regulation.

As used herein, by "modulating" is meant increasing or decreasing cell growth. By "cell growth" is meant any change in cell number or size, including, without
30 limitation, increase or decrease in cell number, increase or decrease in rate of cell division, increase or decrease in rate of cell death, and/or increase or decrease in cell size. Standard methods for measuring cell growth include standard apoptosis assays (e.g., TUNEL assays, DNA fragmentation, trypan blue exclusion) and cell proliferation assays

(*e.g.*, ^3H -thymidine incorporation). It will be appreciated that the degree of modulation of cell growth provided by a plexin polypeptide in a given assay will vary, but one of skill in the art can readily determine the statistically significant change in cell growth of a cell exposed to a plexin polypeptide.

5 By "immune regulation" is meant increasing or decreasing the biological functions of immune cells (*i.e.*, cells involved in an immune response). Immune cells include, without limitation, lymphocytes (T and B), NK cells, dendritic cells, myeloid cells (*e.g.*, macrophages and neutrophils), and other hematopoietic cells involved in an immune response.

10 By "nucleic acid molecule" or "nucleic acid" as used herein, is meant any deoxyribonucleic acid (DNA) or ribonucleic acid (RNA), including, without limitation, complementary DNA (cDNA), genomic DNA, RNA, heteronuclear RNA (hnRNA), messenger RNA (mRNA), DNA/RNA hybrids, or synthetic nucleic acids (*e.g.*, an oligonucleotide) comprising ribonucleic and/or deoxyribonucleic acids or synthetic
15 variants thereof. The nucleic acid of the invention includes, without limitation, an oligonucleotide or a polynucleotide. The nucleic acid can be single stranded, or partially or completely double stranded (duplex). Duplex nucleic acids can be homoduplex or heteroduplex.

By "polypeptide" is meant any molecule comprising two or more amino acids
20 joined together with a peptide bond, regardless of length or post-translational modifications (*e.g.*, without limitation, glycosylation, lipidation, acetylation, or phosphorylation). Useful plexin polypeptides of the invention include, without limitation, the full length plexin polypeptides having the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 or 10; an extracellular domain of the polypeptide having the amino
25 acid sequence 1 to 1199 of SEQ ID NO: 2; 1 to 1099 of SEQ ID NO: 4; 1 to 1270 of SEQ ID NO: 6 or 1 to 199 of SEQ ID NO: 8, with its associated signal peptide; or an extracellular domain of the polypeptide having the amino acid sequence 19 to 1199 of SEQ ID NO: 2; 24 to 1099 of SEQ ID NO: 4; or 43 to 1270 of SEQ ID NO: 6, lacking its associated signal peptide; an intracellular domain of the polypeptide having the
30 amino acid sequence of SEQ ID NOS: 2, 4, 6 or 8; and polypeptides, the amino acid sequence of which comprises about residues 1-18 (putative signal sequence), 19-518 (sema domain), 451-530 (1° MRS), 601-680 (2° MRS), 751-830 (3° MRS), 800-1010 (G-P repeats) or 1196-1215 (putative transmembrane domain) of SEQ ID

NO: 2; about residues 1-23 (putative signal sequence), 24-507 518 (sema domain) or 1100-1119 (putative transmembrane domain) of SEQ ID NO: 4; or about residues 1-42 (putative signal sequence), 43-600 (sema domain), 541-620 (1° MRS), 691-770 (2° MRS), 831-910 (3° MRS), 900-1110 (G-P repeats) or 1270-
5 1293 (putative transmembrane domain) of SEQ ID NO: 6; or about residue 8-49 (1° MRS) or 154-199 (2° MRS) of SEQ ID NO: 8.

By "isolated" is meant a compound (*e.g.*, a nucleic acid molecule or a protein) that has been separated from components (*e.g.*, nucleic acid molecules, proteins, lipids, and/or carbohydrates) which naturally accompany it. Water, buffers, and other small
10 molecules (*e.g.*, molecules having a molecular weight of less than about 1000 daltons) may accompany an isolated compound of the invention. Preferably, an isolated compound is at least 70%, by weight, free from components which naturally accompany it. More preferably, an isolated is at least 75%, by weight, free from components which naturally accompany it; still more preferably, at least 80%, by
15 weight, free; even more preferably, at least 85%, by weight, free; and even more preferably, at least 90%, by weight, free from components which naturally accompany it. Most preferably, a substantially purified compound is at least 95%, by weight, free from components which naturally accompany it.

Where the isolated compound is a nucleic acid molecule, the isolated nucleic acid
20 molecule is separated from other nucleic acids (*e.g.*, genes or transcripts) or proteins which, in the naturally-occurring genome of the organism from which the nucleic acid molecule was derived, flanked the nucleic acid molecule. Isolated nucleic acid molecules therefore include, without limitation, a recombinant nucleic acid molecule incorporated into a plasmid or other vector (*e.g.*, a replication-defective virus); a
25 recombinant nucleic acid molecule incorporated into the genome of a prokaryotic or eukaryotic organism; or a nucleic acid molecule which exists as a separate molecule independent of other nucleic acids (*e.g.*, a PCR product, a chemically synthesized nucleic acid molecule, or a nucleic acid molecule produced by restriction endonuclease digestion). Purification of a nucleic acid molecule can be accomplished and measured by
30 any standard method including, without limitation, sequence analysis, chemical synthesis, PCR, CsCl gradient, phenol:chloroform extraction, ethanol precipitation, Southern or Northern blotting analysis followed by band extraction and purification, and

high performance liquid chromatography (HPLC; see, *e.g.*, Fisher (1980) Laboratory Techniques in Biochemistry and Molecular Biology, Work and Burdon (eds.), Elsevier).

Thus, in one non-limiting example, to obtain an isolated nucleic acid molecule encoding a plexin polypeptide, a nucleic acid molecule is chemically synthesized on a standard oligonucleotide synthesis machine. The resulting single stranded molecule is then subjected to second strand synthesis to form a duplex DNA molecule, which is then ligated into a plasmid capable of replication in a prokaryotic or eukaryotic cell. The nucleic acid molecule is then replicated in the cell, purified (*e.g.*, by CsCl gradient), and subjected to sequence analysis.

In certain embodiments of the first aspect of the invention, the nucleic acid molecule has a nucleic acid sequence comprising SEQ ID NOS: 1, 3, 5, 7 or 9. Preferably, the nucleic acid molecule of the invention has not more than 500 nucleotides flanking each of the 5' and 3' ends of SEQ ID NOS: 1, 3, 5, 7 or 9. In certain embodiments, the plexin polypeptide has an amino acid sequence that comprises SEQ ID NOS: 2, 4, 6, 8 or 10. Preferably, the plexin polypeptide of the invention has not more than 50 amino acid residues flanking each of the N-terminal and C-terminal ends of SEQ ID NOS: 2, 4, 6, 8 or 10.

In certain embodiments of the first aspect of the invention, the nucleic acid molecule hybridizes under stringent conditions (as defined herein) to SEQ ID NOS: 1, 3, 5, 7 or 9.

The invention also includes nucleic acid molecules that hybridize under stringent hybridization conditions (as defined herein) to all or a portion of the nucleotide sequence represented by SEQ ID NOS: 1, 3, 5, 7 or 9 or its complement. The hybridizing portion of the hybridizing nucleic is at least 80%, *e.g.*, at least 95%, or at least 98%, homologous to the sequence of a portion or all of a nucleic acid encoding a polypeptide having the amino acid sequence of SEQ ID NOS: 2, 4, 6, 8 or 10, or its complement. Hybridizing nucleic acids of the type described herein can be used, for example, as a cloning probe, a primer (*e.g.*, a PCR primer) or a diagnostic probe.

Hybridization of the oligonucleotide probe to a nucleic acid sample typically is performed under stringent conditions. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions. If sequences are to be identified that are related and substantially

identical, rather than identical, then it is useful to first establish the lowest temperature at which only homologous hybridization occurs with a particular concentration of salt (*e.g.*, SSC or SSPE). Then, assuming that 1% mismatch results in a 1°C decrease in the T_m , the temperature of the final wash in the hybridization reaction is reduced

5 accordingly (for example, if the sequences have > 95% identity with the probe are sought, the final wash temperature is decreased 5°C). In practice, the change in the T_m can be between 0.5 C and 1.5 C per 1% mismatch. "Stringent conditions" involve hybridization at 68°C in 5x SSC/5x Denhardt's solution/1.0% SDS, and washing in 0.2xSSC/0.1% SDS at room temperature. "Moderately stringent conditions" include

10 washing in 3xSSC at 42°C. The parameters of salt concentration and temperature can be varied to achieve the optimal level of identity between the probe and the target nucleic acid. Additional guidance regarding such conditions is readily available in the art, for example, by Sambrook *et al.*, *supra*; and Ausubel *et al.*, *supra*.

Nucleic acid sequence homology (as well as amino acid sequence homology) can

15 be measured according to standard methods. Unless otherwise specified, as used herein used herein, "percent homology" of two amino acid sequences or of two nucleic acids is determined using the algorithm of Karlin and Altshul (*Proc. Natl. Acad. Sci. USA* **87**: 2264-2268, 1990), modified as in Karlin and Altschul (*Proc. Natl. Acad. Sci. USA* **90**: 5873-5877, 1993). Such an algorithm is incorporated into the NBLAST and XBLAST

20 programs of Altschul *et al.* (*J. Mol. Biol.* **215**: 403-410, 1990). BLAST nucleotide searches are performed with the NBLAST program, e (score) = 100, word length = 12, to obtain nucleotide sequences homologous to a nucleic acid molecule of the invention. BLAST protein searches are performed with the XBLAST program, e (score) = 50, word length = 3, to obtain amino acid sequences homologous to a reference polypeptide (*e.g.*,

25 SEQ ID NO: 2). To obtain gapped alignments for comparison purposes, Gapped BLAST is utilized as described in Altschul *et al.* (*Nucleic Acids Res.* **25**: 3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) are used, namely $e=10$; $w=11$ for nucleic acid; $w=3$ for amino acid (the Blosum 62 scoring matrix); low complexity

30 sequence filtering. The default settings of BLAST emphasize regions of local alignment to detect relationships among sequences which share only isolated regions of similarity (Altschul *et al.*, *J. Mol. Biol.* **215**: 403-410 (1990). See <http://www.ncbi.nlm.nih.gov>.

Thus, in a non-limiting example to obtain an isolated nucleic acid molecule encoding a plexin polypeptide, a nucleic acid molecule having the sequence of SEQ ID NOS: 1, 3, 5 or 7 is used to probe a cDNA library under stringent conditions according to standard techniques (see., *e.g.*, Ausubel *et al.*, *supra*). Upon identification of a positive clone (*i.e.*, a clone that hybridizes to SEQ ID NOS: 1, 3, 5 or 7 under stringent conditions), that clone is expanded and subjected to sequence analysis. A nucleic acid molecule having a nucleic acid sequence that is at least 70% identical, preferably at least 75% identical, more preferably, at least 80% identical, still more preferably at least 85% identical, even more preferably, at least 90% identical, and most preferably at least 95% identical (as measured by the basic BLAST program of NCBI on default settings) to SEQ ID NOS: 1, 3, 5 or 7 is a nucleic acid molecule of the invention.

In a second aspect, the invention provides four novel isolated plexin polypeptides.

"Isolated" is as defined for the first aspect of the invention. Where the isolated compound is a polypeptide, the isolated polypeptide is separated from organic molecules, such as nucleic acid molecules, polypeptides, and/or carbohydrates, which, in the naturally-occurring organism from which the polypeptide was derived, accompany the polypeptide. Isolated polypeptides therefore also include a recombinant polypeptide (*e.g.*, a human polypeptide expressed in an insect cell), or a chemically synthesized polypeptide. Purification of a polypeptide can be accomplished and measured by any standard method including, without limitation, chemical synthesis, recombinant polypeptide expression in prokaryotic or eukaryotic cells, affinity chromatography, Western blotting analysis, SDS-PAGE analysis, and/or HPLC.

In accordance with this aspect, the invention provides all derivatives, mutants, truncations, and/or splice variants of the four novel plexin polypeptides, so long as these derivatives, mutants, truncations, and/or splice variants share at least 60% amino acid sequence homology with SEQ ID NOS: 2, 4, 6 or 8, preferably, at least 65% sequence homology, more preferably, at least 70% sequence homology, more preferably, at least 75% sequence homology, more preferably, at least 80% sequence homology, still more preferably at least 85% sequence homology, even more preferably, at least 90% sequence homology, and most preferably at least 95% sequence homology with SEQ ID NOS: 2, 4, 6 or 8 as determined using the basic BLAST program of the National Center for Biotechnology (NCBI; National Library of Medicine, Bethesda, MD), using the

default settings defined therein using the sequence of the four novel plexin derivative, mutant, truncation and/or splice variance as the probe.

Preferred plexin polypeptide derivatives include polypeptides whose sequences differ from the sequence given in SEQ ID NOS: 2, 4, 6 or 8, by one or more conservative amino acid substitutions, or by one or more non-conservative amino acid substitutions, deletions or insertions which do not abolish the biological activity of the plexins. Conservative amino acid substitutions typically include the substitution of one amino acid for another with similar biochemical characteristics, such as polarity, size, and/or charge. Non-limiting examples of conservative substitutions are substitutions within the following groups: valine, glycine, alanine, valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine, phenylalanine, and tyrosine.

Useful methods for mutagenesis to generate plexin mutants are known in the art (see, e.g., Sambrook *et al.*, *supra*; Ausubel *et al.*, *supra*). Preferred methods for mutagenesis are described in PCT Publication WO99/12965 and include PCR mutagenesis and saturation mutagenesis. A library of random amino acid sequence variants can also be generated by the synthesis of a set of degenerate oligonucleotide sequences.

In certain embodiments of the second aspect of the invention, the plexin polypeptide has a sequence comprising the sequence of SEQ ID NOS: 2, 4, 6 or 8. In one non-limiting example, in accordance with the invention, an isolated plexin polypeptide comprising the sequence of SEQ ID NOS: 2, 4, 6 or 8 can chemically synthesized according to standard techniques (e.g., at a commercial peptide generating facility).

For example, a putative plexin polypeptide is purified and subjected to N-terminal sequencing to determine its amino acid sequence. The amino acid sequence of the polypeptide is then compared to SEQ ID NOS: 2, 4, 6, 8 or 10 (as measured by the basic BLAST program of NCBI on default settings). A polypeptide that shares at least 60% homology with SEQ ID NOS: 2, 4, 6, 8 or 10 is a plexin polypeptide of the invention.

In another example, purification of a plexin polypeptide is facilitated by the addition of a tag to the polypeptide that enables purification of the tagged polypeptide. Non-limiting examples of a tag include a hemagglutinin (HA) tag, a his tag, a GST tag, a

FLAG-tag, and a myc tag. Thus, a nucleic acid molecule of the first aspect is engineered using standard molecular biology techniques to incorporate the nucleic acid sequence encoding the tag. The engineered nucleic acid molecule is then introduced and positioned for expression in an appropriate cell to produce the recombinant tagged polypeptide, which can then be readily purified by binding of the tag to its substrate. For example, the his tag binds to Ni-NTA agarose. Likewise, a GST (glutathione S-transferase) tag binds to glutathione agarose beads. Both his tag and GST tag expression and purification kits are commercially available from PharMingen (San Diego, CA). Likewise, myc-tagged plexin polypeptide are produced by cells introduced with a nucleic acid molecule encoding the tagged protein and positioned for expression in the cell.

It will be appreciated that particularly useful polypeptides of this aspect of the invention are secreted by the cell in which they are produced, thus facilitating purification of the polypeptide from the culture media in which the cells have been maintained, without requiring lysis of the cell.

In a third aspect, the invention provides a cell engineered to comprise a nucleic acid molecule encoding one of the four plexin polypeptides. By "engineered" is meant that the cell of the invention has been modified by standard molecular biology techniques. Where the cell is "engineered to comprise a nucleic acid molecule," standard molecular biology techniques have been employed to introduce the indicated nucleic acid molecule into the cell, either by transformation or transfection of the cell with a plasmid, or by infection or transduction of the cell with a recombinant virus.

The nucleic acid molecule of the first aspect of the invention is preferably subcloned into a plasmid or vector (for example, but not limited to, a vector used to generate a recombinant virus), wherein the nucleic acid molecule is positioned for expression in the plasmid or vector. The plasmid or vector is then introduced into a cell by standard techniques to produce an engineered cell in accordance with the third aspect of the invention.

In certain embodiments of the third aspect, the cell is a prokaryotic cell (*e.g.*, a bacterium). For example, *E. coli* cells (*e.g.*, DH5 α) are transformed (using, *e.g.*, electroporation) with a bacterial plasmid (*i.e.*, a plasmid containing an *E. coli* origin of replication) containing a nucleic acid molecule of the first aspect of the invention. The transformed bacteria are selected using, for example, an antibiotic-resistance encoding nucleic acid molecule (*e.g.*, ampicillin resistance) on the plasmid such that the antibiotic

resistance protein is expressed by the transformed bacteria. The transformed bacteria are then propagated, and can be cryopreserved and stored frozen in glycerol.

Those of skill in the art will appreciate that in accordance with the third aspect of the invention, a nucleic acid molecule encoding one of the four plexin polypeptides may be introduced into a large variety of cells. For example, a nucleic acid molecule encoding one of the four plexin polypeptides can be introduced into prokaryotic cells (*e.g.*, bacteria), and any eukaryotic cell into which an exogenous nucleic acid molecule may be introduced. Thus, in certain embodiments of the third aspect of the invention, the cell is a eukaryotic cell. Eukaryotic cells according to this aspect of the invention that comprise a nucleic acid molecule encoding one of the four plexin polypeptides include, without limitation, yeast cells, plant cells, insect cells, and mammalian cells. Within the category of mammalian cells are cells from any mammalian species (including, without limitation, mouse, hamster, monkey, human), of any lineage (including, without limitation, lymphocyte, fibroblast, stem cell), and may be an immortalized cell, or a non-immortalized cell. Cells, as well as plasmids and/or vectors (*e.g.*, vectors that can be packaged to form infectious virus particles), are commercially available, for example, from the American Type Culture Collection ("ATCC"; Manassas, VA).

In certain embodiments of the third aspect of the invention, the nucleic acid molecule is positioned for expression in the cell. By "positioned for expression" is meant that the nucleic acid molecule is operably linked to at least one regulatory sequence which directs the transcription and translation of the nucleic acid molecule in a cell, such that the cell engineered to comprise the nucleic acid molecule produces (*i.e.*, expresses) the protein encoded by the nucleic acid molecule. By "operably linked" is meant that the nucleic acid molecule and the regulatory sequence are connected in a such a way as to permit expression of the nucleic acid molecule when the nucleic acid molecule is present in a cell. Regulatory sequences include, without limitation, promoters, enhancers, IRES sequences, and polyadenylation signals. Since plexin polypeptides are involved in immune regulation and the modulation of cell growth, it may be desirable to operably link a nucleic acid molecule encoding one of the four plexin polypeptides to an inducible promoter.

The four plexin polypeptides that are encoded by the nucleic acid molecules do not necessarily include the transmembrane domain of the four plexin polypeptides, and so may be produced by the cell as an intracellular polypeptide or a soluble secreted

polypeptide. For example, if the polypeptide fragment is secreted by the cell, it can be purified from the conditioned growth media of the transfected cells, without having to lyse the cells. Likewise, although a soluble intracellular polypeptide fragment is purified from only lysed cells, the fragment, being soluble, does not have to be extracted from the cell membrane; thus, different lysis conditions may be used to obtain purified soluble intracellular polypeptide fragment as compared to the lysis conditions required to obtain purified full length plexin polypeptides (which has a transmembrane domain).

Protein expression systems have been established for a variety of cells and are known to those of skill in the art. Cells are also commercially available from the ATCC, and a variety of protein expression system kits are commercially available from, for example, Invitrogen Corp. (Carlsbad, CA), Clontech Laboratories (Palo Alto, CA), PharMingen (San Diego, CA), Promega Corp. (Madison, WI), and Stratagene (La Jolla, CA).

For example, a nucleic acid molecule encoding one of the four plexin polypeptides is operably linked to bacterial regulatory sequences (e.g., T7 late promoter or bacteriophage regulatory sequences), and the resulting nucleic acid molecule is used to transform bacterial cells, where the transformed bacterial cells produce one of the four plexin polypeptides. In another example, a nucleic acid molecule encoding one of the four plexin polypeptides is operably linked to baculovirus regulatory sequences in a baculovirus vector. Recombinant baculovirus are then generated and used to transduce insect cells (using, for example, the expression kit commercially available from Clontech Laboratories). The transduced insect cells comprise a nucleic acid molecule encoding one of the four plexin polypeptides positioned for expression in the insect cell, and thus produce one of the four plexin polypeptides.

Mammalian cells are widely used as protein expression systems. For example, a mammalian cell may be transduced with a recombinant retrovirus or adenovirus comprising a nucleic acid molecule encoding one of the four plexin polypeptides operably linked to regulatory sequences that are either endogenous to the particular virus or exogenous to the virus (e.g., a CMV promoter in a retroviral vector). The transduced mammalian cell is then propagated *in vitro* in tissue culture, *in vivo* (e.g., in an immunocompromised animal), and/or cryopreserved and stored frozen in DMSO.

In another example, mammalian cells are transfected with an expression plasmid comprising a nucleic acid molecule encoding one of the four plexin polypeptides

operably linked to one or more regulatory sequences on the plasmid. By "expression plasmid" is meant a plasmid in which an inserted nucleic acid molecule of interest (*e.g.*, encoding one of the four plexin polypeptides, a plexin chimeric molecule, or tagged plexin polypeptide) is operably linked to at least one regulatory sequence such that when
5 the expression plasmid containing the inserted nucleic acid molecule of interest is introduced (*e.g.*, by transfection) into a cell, the inserted nucleic acid molecule is positioned for expression in that cell. The nucleic acid molecule in the expression plasmid, upon being introduced into the cell, is thus positioned for expression in that cell, and enables the cell to produce one of the four plexin polypeptides encoded by the
10 nucleic acid molecule.

In one non-limiting example, a nucleic acid molecule according to the first aspect of the invention is inserted into a standard mammalian expression plasmid (*e.g.*, pCDNA3.1 commercially available from Invitrogen Corp., Carlsbad, California), such that the inserted nucleic acid molecule encoding one of the four plexin polypeptides is
15 operably linked to the regulatory sequences in the mammalian expression plasmid. Mammalian cells are then transfected with this expression plasmid (using, *e.g.*, CaPO₄ or DEAE-dextran). Where the expression plasmid contains an antibiotic-resistance encoding nucleic acid molecule (*e.g.*, neomycin resistance on the pCDNA3.1 plasmid) such that the antibiotic resistance protein is expressed by the transfected cells, transfected
20 cells may be selected for the ability to grow in the presence of the antibiotic. The transfected cells may then be propagated and cryopreserved and stored in frozen in DMSO.

In a fourth aspect, the invention provides an isolated nucleic acid molecule encoding a chimeric molecule comprising at least two segments, wherein one of the
25 segments comprises one of the four plexin polypeptides. By "chimeric molecule" is meant a protein that comprises at least two segments of polypeptide joined together by any means, including, without limitation, a covalent bond (*e.g.*, peptide bond), a non-covalent bond (*e.g.*, ionic bond or hydrogen bond) or by a chemical crosslinker. It should be noted that one of the four plexin polypeptides that has been tagged is within the
30 definition of a chimeric molecule.

In certain embodiments of the fourth aspect of the invention, the nucleic acid molecule encoding the segment of a chimeric molecule comprising one of the four plexin

polypeptides hybridizes under stringent conditions to SEQ ID NO: 1, 3, 5 or 7.

"Stringent conditions" are as described above for the first aspect of the invention.

Standard molecular biology techniques may be employed to generate nucleic acid molecules encoding chimeric molecules according to the fourth aspect of the invention.

- 5 For example, a nucleic acid molecule encoding the extracellular domain of one of the four plexin polypeptides may be joined, in frame, to a nucleic acid molecule encoding the constant region of an immunoglobulin molecule (see, *e.g.*, Chamow S.M., Antibody Fusion Proteins, John Wiley & Sons, New York, 1999). By "in frame" is meant that a first nucleic acid molecule is ligated to a second nucleic acid molecule such that the each
10 of the amino acid sequences of the polypeptides encoded by each of the first and the second nucleic acid molecules is not frame-shifted.

- In one non-limiting example, a chimeric molecule comprising the extracellular domain of one of the four plexin polypeptides including the amino acid sequence of SEQ ID NOS: 2, 4, 6 or 8 is generated. In this example, a nucleic acid molecule encodes
15 amino acid residue number 1(19) through about amino acid residue number 1199 of SEQ ID NO: 2; amino acid residue number 1(24) through about amino acid residue number 1099 of SEQ ID NO: 4; amino acid residue number 1(43) through about amino acid residue number 1270 of SEQ ID NO: 6 and amino acid residue number 1 through about amino acid residue number 199 of SEQ ID NO: 8 with its associated signal peptide
20 (parenthesis depicts about the beginning of the amino acid sequence of the extracellular domain lacking its signal peptide). This nucleic acid molecule is fused in frame with a nucleic acid molecule encoding the constant region of an immunoglobulin, such that the chimeric molecule encoded by the resulting nucleic acid molecule generally has the following structure:

25

N-terminus	extracellular domain of SEQ ID NO: 2 with or lacking its signal peptide	amino acids from the constant region of an Ig molecule	C-terminus
N-terminus	extracellular domain of SEQ ID NO: 4 with or lacking its signal peptide	amino acids from the constant region of an Ig molecule	C-terminus
N-terminus	extracellular domain of SEQ ID NO: 6 with or lacking its signal peptide	amino acids from the constant region of an Ig molecule	C-terminus
N-terminus	extracellular domain of SEQ ID NO: 8 with or lacking its signal	amino acids from the constant region of an Ig molecule	C-terminus

	peptide		
--	---------	--	--

The heavy chain class (e.g., IgG, IgA, IgM, IgD, or IgE) can be varied in this chimeric molecule depending upon which constant region is used. Nucleic acid molecules encoding the constant region of various immunoglobulin (Ig) heavy chains are known
5 (see, e.g., Zettlmeissl et al., *DNA Cell Biol.* **9**(5):347-53, 1990) Indeed, expression plasmids are available, into which the nucleic acid molecule of interest (i.e., a nucleic acid molecule encoding an extracellular domain of the polypeptide of SEQ ID NO: 2; SEQ ID NO: 4; SEQ ID NO: 6; or SEQ ID NO: 8) can be inserted, and the resulting plasmid introduced into a cell to produce one of the four extracellular plexin-Ig chimeric
10 molecule s (see, e.g., Zettlmeissl et al., *supra*; Miller et al., *J. Exp. Med.* **178** (1): 211-222, 1993).

Any variety of chimeric molecule carrying the extracellular domains of one of the four plexin polypeptide may be generated. For example, the extracellular domain of one of the four plexin polypeptides can be myc-tagged, his-tagged, or FLAG tagged
15 according to standard molecular biology techniques.

Such extracellular proteins are particularly useful for identifying ligands to which the extracellular domain of one of the four plexin polypeptides bind. For example, extracellular plexin-D1-Ig chimera can be immobilized on a protein A-sepharose column. Molecules suspected of binding the extracellular domain of plexin-D1 are added to the
20 column, to which the molecule that binds to the extracellular domain of plexin-D1 adhere, and the non-binding molecules flow through the column. The extracellular plexin-D1-binding molecules are readily eluted, for example, by changing the pH or ion concentration of the elution buffer.

Extracellular plexin proteins are also used to identify cells expressing the ligand
25 of plexin extracellular domain on their cell surface (and thereby also identify the ligand itself). For example, cells are incubated with a FLAG-tagged plexin extracellular domain chimeric molecule. A FLAG-tagged plexin extracellular domain chimeric molecule is generated. An anti-FLAG antibody that is detectably labeled is then added to the cells. By "detectably labeled" is meant any means for marking and identifying the presence of a
30 molecule. Detectable labels include, without limitation, radioactive labels (e.g., ^{32}P or ^{35}S) and fluorophore labels (e.g., FITC, phycoerythrin, or rhodamine). For example, FITC-labeled anti-FLAG antibodies are commercially available from Babco, Richmond,

CA. The "stained" cells (*i.e.*, cells incubated first with the FLAG-tagged plexin extracellular domain chimeric molecule and then with the FITC-labeled anti-FLAG antibody), are then subjected to flow cytometry analysis to select those cells that are labeled with FITC, and so express a molecule that binds to the extracellular domain of one of the four plexin polypeptides. The FITC labeled cells are then further manipulated (*e.g.*, characterized to determine which cells express the plexin polypeptide ligand).

The ligand of the plexin extracellular domain is itself identified, for example, by lysing the cells, adding the lysate to one of the four plexin extracellular domain-Ig chimeric molecule columns described above, and purifying the ligand. The ligand is then sequenced by N-terminal sequencing.

In another non-limiting example, the intracellular domain of one of the four plexin polypeptides is used as "bait" in a yeast two-hybrid system to identify ligands that interact with the intracellular domain of one of the four plexins described herein. For general description of the two-hybrid system, see U.S. Patent Nos. 5,283,173; 5,468,614; and 5,695,941. In this example, a nucleic acid molecule encoding from about amino acid residue number 143 through at least amino acid residue number 214 of SEQ ID NO: 2 is inserted into a standard DNA binding domain expression plasmid (*e.g.*, the GAL4 DNA binding domain plasmid in the Interactor kit commercially available from PharMingen (San Diego, CA). (It will be understood that the nucleic molecule may encode amino acid residue number 138-148 through at least amino acid residue number 214 of SEQ ID NO: 2.) A variety of cDNA libraries in transcriptional activation domain vectors are available (*e.g.*, from Clontech, Palo Alto CA). The cDNA libraries are screened employing standard methods (see, *e.g.*, the methods employed in U.S. Patent No. 5,780,262) to identify cDNA clones encoding a ligand that binds to the intracellular domain of one of the four plexin polypeptides. One preferable cDNA library screened in this example is a cDNA library generated from an immune cell (*e.g.*, a lymphocyte or NK cell).

In a fifth aspect, the invention provides a purified chimeric molecule comprising one of the four plexin polypeptides. Methods for purifying proteins are as described for the second aspect of the invention.

In a sixth aspect, the invention provides a cell engineered to comprise a nucleic acid molecule encoding a chimeric molecule comprising at least two segments, wherein one of the segments comprises one of the four plexin polypeptides. As described for the

third aspect of the invention, a nucleic acid encoding a chimeric molecule comprising one of the four plexin polypeptides may be introduced into any variety of cells. In certain embodiments, the cell is a prokaryotic cell or a eukaryotic cell. In certain embodiments, the eukaryotic cell is a yeast cell or a mammalian cell (*e.g.*, a human cell).

5 In a seventh aspect, the invention provides an isolated binding agent that specifically binds one of the four plexin polypeptides, or specifically binds a chimeric molecule comprising a segment comprising one of the four plexin polypeptides. In certain embodiments, the plexin protein has an amino acid sequence comprising SEQ ID NOS:2, 4, 6 or 8.

10 By "specifically binds" is meant a binding agent (*e.g.*, an antibody) that binds to its specific target (*e.g.*, one of the four plexin polypeptides) with greater affinity than it binds to other molecules. Preferably, where the binding agent is an antibody, the antibody preferably specifically binds to its specific target with a dissociation constant (K_D) of at least 10^{-5} M, more preferably, 10^{-6} M, even more preferably 10^{-7} M, and most
15 preferably, the binding agent specifically binds to its specific target with a K_D of at least 10^{-8} M.

Preferably, the binding agent of this aspect of the invention is an antibody, such as a monoclonal antibody or a polyclonal antibody, or a fragment of an antibody that specifically binds one of the four plexin polypeptides. Standard methods may be
20 employed to generate both monoclonal and polyclonal antibodies that specifically bind to one of the four plexin polypeptides of the invention. See, *e.g.*, Ausubel et al., *supra*; Coligan, J.E. et al., Current Protocols in Immunology, John Wiley & Sons, New York (1991); and Delves, P.J., Antibody Production: Essential Techniques, John Wiley & Sons, New York (1997). Briefly, the plexin polypeptides of the present invention,
25 purified according to the methods described for the second aspect of the invention, are used to immunize rabbits (*e.g.*, for polyclonal antibodies) or mice (*e.g.*, for monoclonal antibodies) to generate antibody-mediated immunity to the four plexin polypeptides used to immunize the animal. For monoclonal antibodies, antibodies can be screened by, *e.g.*, ELISA, to identify those antibodies that show the highest affinity for the immunizing
30 plexin protein of polypeptide fragment. The cloned cell producing the high affinity monoclonal antibody can then propagated *in vitro* (where the antibody is purified from the culture supernatant) or *in vivo* (where the antibody is purified from ascites fluid), and

can also be cryopreserved and stored frozen at, *e.g.*, -70°C in DMSO, to provide a potentially limitless supply of monoclonal antibody.

In addition to intact monoclonal and polyclonal antibodies, the invention also provides various antibody fragments, such as Fab, F(ab')₂, Fv, and sFv fragments.

- 5 Recombinant, chimeric, and humanized antibodies are also provided.

Recombinant "humanized antibodies" which specifically bind to one of the four plexin polypeptides can be synthesized according to methods known in the art (see, *e.g.*, Green L.L. *et al.*, *Nature Genetics* 7: 13-21, 1994 for fully humanized antibodies expressed in transgenic animals; see also U.S. Patent Nos. 5,693,761; 5,777,085; and 10 5,585,089). Humanized antibodies are chimeras comprising mostly human IgG sequences into which at least portions of the regions responsible for specific antigen-binding (*e.g.*, CDR's) have been inserted. Animals are immunized with the desired antigen, the corresponding antibodies are isolated, and the portion of the variable region sequences responsible for specific antigen binding are removed. The animal-derived 15 antigen binding regions are then cloned into the appropriate position of human antibody genes in which the antigen binding regions have been deleted. Humanized antibodies minimize the use of heterologous (*i.e.*, inter-species) sequences in human antibodies, and thus are less likely to elicit immune responses in the treated subject (see also, *e.g.*, U.S. Patent No. 5,807,715).

- 20 Construction of different classes of recombinant antibodies can also be accomplished by making chimeric or humanized antibodies comprising nonhuman variable domains and human constant domains (CH1, CH2, CH3) isolated from different classes of immunoglobulins. For example, antibodies with increased antigen binding site valencies can be recombinantly produced by cloning the antigen binding 25 site into vectors carrying the human chain constant regions (see, *e.g.*, Arulanandam *et al.*, *J. Exp. Med.* 177: 1439-1450, 1993).

In addition, standard recombinant DNA techniques can be used to alter the binding affinities of recombinant antibodies with their antigens by altering amino acid residues in the vicinity of the antigen binding sites. The antigen binding affinity of a 30 humanized antibody can be increased by mutagenesis based on molecular modeling (see, *e.g.*, Queen *et al.*, *Proc. Natl. Acad. Sci.* 86: 10029-10033, 1989).

Also provided in the invention are plexin polypeptide-specific single polypeptide chain antibodies (see general methods in U.S. Patent Nos. 4,946,788 and

4,704,692); single domain antibodies (Ward, E.S. et al., *Nature* **341**: 544-546, 1989); and chimeric antibodies (U.S. Patent No. 4,816,567).

Binding agents that specifically bind the plexin polypeptides of the present invention are useful, for example, in determining expression levels of the plexin polypeptides in various tissues of the body, Western blotting analysis, and immunochromatography. Particularly, binding agents that specifically bind the plexin polypeptides are useful for binding the plexin polypeptide on a cell expressing the plexin polypeptide, thereby activating the cell.

A binding agent that specifically binds one of the four plexin polypeptides, for example, is effective as an immune modulator. Additional applications include, without limitation, an injectable formulation comprising a binding agent that specifically binds one of the four plexin polypeptides that is useful to antagonize activity in a disease involving aberrant immune regulation or a disease involving aberrant cell growth.

In an eighth aspect, the invention provides an isolated antisense oligonucleotide complementary to a portion of a nucleic acid molecule encoding one of the four plexin polypeptides. In certain embodiments, hybridization of the antisense oligonucleotide to the nucleic acid molecule inhibits transcription or translation of the nucleic acid molecule.

By two nucleic acid molecules being "complementary" to one another is meant that the first nucleic acid molecule (*e.g.*, an oligonucleotide) is able to form Watson-Crick base pair hydrogen bonds (*i.e.*, hybridize) with the second nucleic acid molecule to form a duplex. The first nucleic acid molecule is thus a "complement" of the second nucleic acid molecule.

The antisense oligonucleotides according to the invention are complementary to a region of a nucleic acid molecule (or a region at the intron/exon boundary of DNA or RNA) that encodes one of the four plexin polypeptides. Preparation of antisense oligonucleotides is well known (see, *e.g.*, Agrawal *et al.*, *Trends Biotechnol.* **10**:152-158, 1992; U.S. Patent No. 5,149,798; Agrawal *et al.*, *Proc. Natl. Acad. Sci. USA* **85**:7079-7083, 1988; Froehler, *Tetrahedron Lett.* **27**:5575-5578, 1986; and Bergot *et al.*, *J. Chromatog.* **559**:35-42, 1992).

In a ninth aspect, the invention provides a method for identifying a nucleic acid molecule encoding one of the four plexin polypeptides, comprising contacting a pool of candidate nucleic acid molecules with a nucleic acid molecule encoding one of the four

plexin polypeptides, wherein hybridization of the nucleic acid molecule encoding one of the four plexin polypeptides under stringent conditions to a candidate nucleic acid molecule identifies the candidate nucleic acid molecule as a nucleic acid molecule that encodes one of the four plexin polypeptides. According to this aspect of the invention, "hybridization" and "stringent conditions" are as defined above for the first aspect of the invention. In certain embodiments, the nucleic acid molecule encoding one of the four plexin polypeptides has a nucleic acid sequence comprising SEQ ID NOS: 1, 3, 5 or 7.

It will be understood that the isolated plexin polypeptides according to the second aspect of the invention, the plexin chimeric molecules according to the fifth aspect of the invention, and binding agents that specifically bind the plexin polypeptides according to the seventh aspect of the invention, are useful as therapeutics to treat an individual suffering from, or suspected of having, a disease or disorder involving aberrant immune regulation or an individual suffering from, or suspected of having, a disease or disorder involving aberrant cell growth, particularly nerve cell growth.

By "disease or disorder involving aberrant immune regulation" is meant any disease or disorder in which an abnormal immune response is generated in response to either self or foreign antigens. Thus, this definition includes, without limitation, autoimmune diseases (*e.g.*, lupus, inflammatory bowel disease, or Diabetes Type 1) and immunosuppressive diseases (*e.g.*, multiple sclerosis or rheumatoid arthritis).

By "disease or disorder involving aberrant cell growth" is meant any disease or disorder in which an abnormal amount of cell growth is observed. "Cell growth" is defined above. Thus, diseases and disorders involving aberrant cell growth include hyperplasia, neoplasia, and cancer, as well as degenerative diseases, such as neurodegenerative diseases.

Preferable therapeutically useful plexin polypeptides are soluble polypeptides (*e.g.*, lacking the hydrophobic transmembrane domain of the plexin polypeptides), particularly soluble polypeptide fragments that are secreted by the cell in which the fragment was produced. In a preferred embodiment the soluble plexin polypeptides are selected from the group consisting of plexin-A-1 (Maestrini et al. 1996 *supra*), plexin-A-2 (Maestrini et al. 1996 *supra*), plexin-A-3 (Maestrini et al. 1996 *supra*), plexin-A-4, plexin-B-1 (Maestrini et al. 1996, *supra*), plexin-B-2, plexin-B-3, plexin-C1 (Comeau et al. 1998 *supra*), plexin-D-1.

In a tenth aspect, the invention provides a method for diagnosing a disease involving aberrant immune regulation or a disease involving aberrant cell growth, comprising comparing the amino acid sequence of one of the four plexin polypeptides from an individual suspected of having the disease with the amino acid sequence of one of the four plexin polypeptides from an unaffected individual, wherein the presence of a difference between the two amino acid sequences identifies the individual suspected of having the disease as having the disease. "Disease or disorder involving aberrant immune regulation" and "disease or disorder involving aberrant cell growth" are as defined above.

By "difference" in the amino acid sequence of one of the four plexin polypeptides from an individual suspected of having the disease or disorder as compared with the amino acid sequence of one of the four plexin polypeptides from an unaffected individual, is meant any mutation that changes the amino acid sequence including substitution, deletion, of addition of one or more amino acid residues.

Thus, in one nonlimiting example, one of the four plexin polypeptides is extracted from cells of an individual suspected of having a disease involving aberrant immune regulation (*e.g.*, using an antibody according to the seventh aspect of the invention). The amino acid sequence of the plexin polypeptide is determined by N-terminal sequencing and compared to the amino acid sequence of one of the four plexin polypeptides from an unaffected individual (*i.e.*, a normal individual of the same species that does not have a disease involving aberrant immune regulation or a disease involving aberrant cell growth). If there is a difference in the two amino acid sequences, the individual suspected of having a disease involving aberrant immune regulation is identified as having a disease involving aberrant immune regulation, and may be treated accordingly.

In certain embodiments of the tenth aspect, the amino acid sequence of the plexin polypeptide from the unaffected individual comprises the sequence of SEQ ID NO: 2, 4, 6, 8 or 10.

The following examples are intended to further illustrate certain preferred embodiments of the invention and are not limiting in nature. Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such

equivalents are considered to be within the scope of this invention, and are covered by the following claims.

Practice of the present invention will employ, unless indicated otherwise, conventional techniques of cell biology, cell culture, molecular biology, microbiology, recombinant DNA, protein chemistry, and immunology, which are within the skill of the art. Such techniques are described in the literature. See, for example, **Molecular Cloning: A Laboratory Manual**, 2nd edition. (Sambrook, Fritsch and Maniatis, eds.), Cold Spring Harbor Laboratory Press, 1989; **DNA Cloning**, Volumes I and II (D.N. Glover, ed), 1985; **Oligonucleotide Synthesis**, (M.J. Gait, ed.), 1984; U.S. Patent No. 4,683,195 (Mullis et al.); **Nucleic Acid Hybridization** (B.D. Hames and S.J. Higgins, eds.), 1984; **Transcription and Translation** (B.D. Hames and S.J. Higgins, eds.), 1984; **Culture of Animal Cells** (R.I. Freshney, ed). Alan R. Liss, Inc., 1987; **Immobilized Cells and Enzymes**, IRL Press, 1986; **A Practical Guide to Molecular Cloning** (B. Perbal), 1984; **Methods in Enzymology**, Volumes 154 and 155 (Wu et al., eds), Academic Press, New York; **Gene Transfer Vectors for Mammalian Cells** (J.H. Miller and M.P. Calos, eds.), 1987, Cold Spring Harbor Laboratory; **Immunochemical Methods in Cell and Molecular Biology** (Mayer and Walker, eds.), Academic Press, London, 1987; **Handbook of Experiment Immunology**, Volumes I-IV (D.M. Weir and C.C. Blackwell, eds.), 1986; **Manipulating the Mouse Embryo**, Cold Spring Harbor Laboratory Press, 1986.

The following Examples are provided to illustrate the present invention, and should not be construed as limiting thereof.

EXAMPLES

Example 1

Identification and cDNA cloning of *plexins* and sequence analysis

Since the coding sequences of human *plexin-B1(SEP)*, *plexin-A2(OCT)* and
5 *plexin-A1(NOV)* were incomplete, we obtained the missing cDNA by RT-PCR; primers
were designed by homology to orthologous murine sequences and corresponding ESTs.
Updated database entries are X87904, X87831 and X87832, respectively. Partial cDNA
of *plexin-A4* was obtained by assembling five overlapping human ESTs (HGI THC
Report: THC203425), deriving from chromosome 7 specific cDNA pools. Another EST
10 from chr. 7 (clone 7B19F10) encodes the cytoplasmic domain of a plexin and
presumably derives from the same gene as *plexin-A4*. *Plexin-B2* cDNA was amplified
by RT-PCR starting from the partial cDNA sequences of clones *MM1* (Shinoura, N.,
Shamraj, O.I., Hugenholz, H., Zhu, J.G., McBlack, P., Warnick, R., Tew, J.J., Wani,
M.A., and Menon, A.G. (1995). Identification and partial sequence of a cDNA that is
15 differentially expressed in human brain tumors. *Cancer Lett* 89, 215-221) and
KIAA0315 (Genbank database); the genomic locus of *SEP-B* was identified due to its
100% sequence identity with clone C22_311 from human chromosome 22. *Plexin-B3*
coding sequence was identified in the genomic sequence of ALD locus on human
chromosome Xq28, using the algorithms HEXON and GENIE. *Plexin-D1* was similarly
20 found in the genomic sequence of chromosome 3 (pac pDJ70i11). The genomic
sequence of *plexin-B1(SEP)*, in the region of the alternative splicing of the extracellular
domain, was obtained using the following primers: sense
5'GCAGCACCTGTGCACCCACAAGGC3' and antisense:
5'TGCAGGCTGGACGGGAGGATGAGG3'. The common donor site is
25 CCATCAG/gtgattgt (position 2028 from ATG); the alternative splice acceptor sites are:
(i) ccccttcag/AGCCC, leading to the canonical plexin-B1 sequence, and (ii)
ctctctcag/GTGAT, leading to "plexin-B1 truncated" variant. All these new sequences
were analyzed using the algorithms BLAST2, NETPHOS (phosphorylation prediction
sites, by Nicolaj Blom), PH-PREDICT and Consensus Protein Secondary Structure
30 prediction at IBCP. The phylogenetic tree was generated using AIIAII algorithm of the
Darwin sequence analysis system (at CBRG).

Example 2

Plexin cDNA expression constructs and protein analysis

Cell transfections were carried out by Calcium phosphate and DEAE-dextran methods, using 5-10 µg of each cDNA (1-2 µg each in case of cotransfections). For transient transfections in COS and BOSC-23 cells the cDNA was cloned in pCDNA3 or derived expression plasmids (Invitrogen). MDCK stable transfectants for *plexin-A3* were obtained using pCEP4 expression plasmid (Invitrogen); the selection was done in the presence of Hygromycin-B (100-200 µg/ml). *Plexin-A3* positive clones of MDCK cells were isolated from two independent transfections, and showed identical biological properties. *Plexin* and neuropilin expression constructs included a VSV- and myc-tag at the N' and C' protein termini, respectively, detected by monoclonal antibodies anti-VSV-G (cat. V-5507, Sigma) and anti-cMyc-tag (cat. OP10-100UG, Calbiochem). "Plexin-B1 truncated" splice variant was expressed from a cDNA fragment isolated by RT-PCR and VSV-tagged at the N' terminus: the encoded amino acid sequence spans up to aa 676 (including the *sema* domain and two MRS motifs). "Plexin-B1-sema" derives from a further deletion of the *plexin-B1* extracellular domain, and exclusively includes the *sema* domain. "Plexin-B1-Δsema" protein mutant includes only the C' terminal half of *plexin-B1* extracellular domain, starting from amino acid 606, i.e. excluding *sema* domain and first MRS but including second and third MRS, transmembrane and intracellular domains.

For immunoprecipitations, cells were lysed with EB buffer (20 mM Tris-HCl pH 7.4, 5 mM EDTA, 150 mM NaCl, 10% glycerol, 1% Triton X-100), in the presence of a cocktail of protease inhibitors and 1mM Na-ortovanadate. Immunoprecipitations were performed at 4°C for 4h with the appropriate antibodies; high stringency washes were performed, in the presence of 1 M LiCl.

For *in vitro* kinase assays, immunopurified proteins were incubated with kinase buffer (50 mM Hepes, 100 µM DTT, 5 mM MnCl₂, 5 mM MgCl₂) in the presence of redivue 5 µCi [γ-³²P] ATP (Amersham) for 10 minutes at 4°C in agitation. Samples were then submitted to SDS-PAGE and autoradiography, or analysed using a Phosphor-Imager system (Molecular Dynamics). Alkali treatment of the polyacrilamide gels was performed with 1M KOH for two hours at 55°C.

Western blots were performed according to standard methods. Specific detection of phospho-tyrosines was done with PY20 MoAb (Trasduction laboratories). Final detection was done with ECL system (Amersham).

Example 3

Semaphorin-SEAP binding assays

Soluble forms of Semaphorin extracellular domains were expressed as chimeric molecules with placental Secreted Alkaline Phosphatase (SEAP) and harvested from the conditioned media of transiently transfected COS or BOSC-23 cells. Serum-free media were concentrated over 100 times using Centricon Plus-20 filters (Millipore) with a molecular weight cutoff of 100 kDa. The AP activity of these media was assessed as described (Flanagan, J.G. and Leder, P. (1990). The kit ligand: a cell surface molecule altered in steel mutant fibroblasts. *Cell* 63, 185-194); the specific activity of chimeric molecules was approx. 1000 U/mg. Concentrated Semaphorin-SEAP were diluted as appropriate in a Hepes buffered saline, additioned with 0.2% BSA, 0.1% NaN₃, 5 mM CaCl₂ and 1 mM MgCl₂ (HBSBA). For binding assays, COS cells transiently transfected with plexins were seeded on 48 well plates to reach confluence, and then incubated with Semaphorin-SEAP preparations (approx 1-5 nM) for 90 minutes at room temperature. The binding was detected as described (Flanagan and Leder, 1990). Binding experiments with plexin-C1/VESPR were as described (Comeau, M.R., Johnson, R., DuBose, R.F., Petersen, M., Gearing, P., VandenBos, T., Park, L., Farrah, T., Buller, R.M., Cohen, J.I., Strockbine, L.D., Rauch, C., and Spriggs, M.K. (1998). A poxvirus-encoded semaphorin induces cytokine production from monocytes and binds to a novel cellular semaphorin receptor, VESPR. *Immunity*, 8, 473-482; He, Z. and Tessier-Lavigne, M. (1997). Neuropilin is a receptor for the axonal chemorepellent Semaphorin III. *Cell* 90, 739-751).

For *in vitro* binding assays, plexin-B1 was purified from cell extracts by immunoprecipitation with anti-VSV antibody. Extracts of mock-transfected cells were used as control samples. After washing, the immunocomplexes were incubated with serial dilutions of CD100-SEAP (prepared as above) for 2 hours at 4°C, in continuous agitation. Samples were then washed 3 times with HBSBA and the bound alkaline phosphatase activity was measured by colorimetric assay using p-nitro-phenyl-phosphate, as described (Flanagan and Leder, 1990). Scatchard analysis was done using Equilibrate (by GertJan C. Veenstra).

Example 4

In situ hybridization analysis

RNA *in situ* hybridization was performed essentially as described (He, Z. and Tessier-Lavigne, M. (1997). Neuropilin is a receptor for the axonal chemorepellent Semaphorin III. *Cell* 90, 739-751). Briefly cDNA fragments of plexin-A1, -A2, and -A3 were used to generate ³⁵S-labeled antisense and sense RNA probes, which were used for *in situ* hybridization histochemistry of cryostat sections of rat embryos.

Example 5

Xenopus turning assay

The methods for injecting mRNA encoding various constructs, and for studying the turning responses of the neurons, are exactly as described previously (Hong, K., Hinck, L., Nishiyama, M., Poo, M.M., Tessier-Lavigne, M., and Stein, E. (1999). A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. *Cell* 97, 927-941); Song, H., Ming, G., He, Z., Lehmann, M., Tessier-Lavigne, M., and Poo, M. (1998). Conversion of neuronal growth cone responses from repulsion to attraction by cyclic nucleotides [see comments]. *Science* 281, 1515-1518).

Example 6

Mixed-culture assays and time-lapse videomicroscopy

Mock-transfected and plexin-A3 overexpressing MDCK cells were seeded with mesenchymal cells (NIH 3T3, KJ29, D17, among others), in multiwell culture plates by 1:4 or 1:1 ratio. NIH and KJ-29 cells were sometimes labeled by addition of DiI (1,1'-diiododecyl-3,3',3'-tetramethylindocarbocyanine perchlorate, Fluka) in the culture medium, 4 hours before harvesting for the assay; clusters of cells marked with this dye are marked in blue (in light microscopy) and emit red epifluorescence (TRITC filter). The repelling effect was observed 16-30 hours after confluency, by contrast phase microscopy using Leica DM IL. The progress of the assays was also monitored by time-lapse video-microscopy (320 minutes recording were converted into 1 minute play). To determine the time-length of cell contacts, for each assay, randomly chosen fibroblasts were followed during several hours and the duration of each contact between their lamellipodia and MDCK cells was measured. The doubling time of cells and their viability during the assay could also be analyzed, and no differences were observed in presence of control or plexin-A3 expressing cells. Substrate adhesion of plexin-A3 overexpressing MDCKs was analyzed by counting attached cells after 30 minutes from

seeding on micro-wells coated with fibronectin, collagen or polylysine, in the absence of calf serum: no differences versus control cells were observed.

Example 7

Apoptosis detection

5 TUNEL reaction (Boehringer detection kit) was performed on mixed cultures of MDCK and NIH3T3 cells, 24 hours after seeding in a 24-well culture plate. The labeling was converted into a colorimetric signal for analysis by light microscopy using the TUNEL-AP detection kit (Boehringer). As a positive control for the induction of apoptosis, the same cells were treated with UV-C (50 $\mu\text{J}/\text{cm}^2$) or 1 μM staurosporin.

10 Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be apparent to those skilled in the art that certain changes and modifications will be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention, which is delineated by the appended claims.

15 Example 8

Plexins are specific receptors for cell surface semaphorins in vertebrates

Plexin-C1 (VESPR) has been shown to bind the soluble viral semaphorins Sema VA and VB (Comeau et al., 1998 *supra*), and we recently found that Drosophila Plexin A (D Plex A) interacts with transmembrane Sema 1a (Winberg et al., 1998 *supra*). We
20 therefore examined in vertebrates whether the extracellular domain of several different cellular semaphorins -fused to alkaline phosphatase- could bind members of the human plexin-A, -B and -C subfamilies. Multiple secreted semaphorins of class 3 (Sema3A, Sema 3C or Sema3F; see below) did not interact with plexins-A1, -A2, -A3, -B1, B2, or -C1 (data not shown). In contrast, plexin-C1(Vespr) specifically bound Sema7A(Sema-
25 K1) (Fig. 2a), a GPI-membrane linked semaphorin (class 7). This result is not entirely unexpected, since Sema7A may represent the cellular counterpart of viral semaphorin SemaVB, previously shown to interact with this plexin (Comeau et al., 1998 *supra*). More interestingly, the class 4 transmembrane semaphorin Sema4D (CD100) did
30 interact strongly and specifically with plexin-B1 (Fig. 2a). Thus the prototypes of two distinct plexin families are the receptors for members of two distinct semaphorin sub-classes. We also found that Sema7A and Sema4D do not bind to neuropilin-1 or -2 alone, nor did co-transfection of either neuropilin with plexin-B1 significantly modify

its binding efficiency (not shown). Neuropilins thus seem so far to function as receptors only for vertebrate semaphorins of class 3.

The affinity constant of Sema4D for plexin-B1 was estimated by Scatchard plot to be in the subnanomolar range ($K_D = 0.9$ nM, Fig. 2b; the estimated K_D of Sema7A for plexin-C1 is 2.1 nM, not shown). These values are consistent with those observed for semaphorins-neuropilins, and fly semaphorin1-Plexin A interactions (He and Tessier-Lavigne, 1997 *supra*; Winberg et al., 1998).

We used two deletion constructs of plexin-B1 to explore the semaphorin binding sites of plexins. Neither the N-terminal half of plexin-B1 extracellular domain ("plexin-B1 truncated", see previous paragraph), nor its C-terminal half ("plexin-B1- Δ sema", see Experimental Procedures) was sufficient alone to bind CD100 (see Fig. 2a), suggesting that the binding of Sema4D depends on multiple structural determinants of the extracellular domain of plexin-B1.

Example 9

Plexins associate with class 3 Semaphorin receptors. Neuropilins

As outlined above, secreted semaphorins of subclass 3 are known to bind neuropilins (He and Tessier-Lavigne, 1997 *supra*; Kolodkin et al., 1997 *supra*; Chen, H., Chedotal, A., He, Z., Goodman, C.S., and Tessier-Lavigne, M. (1997) "Neuropilin-2, a novel member of the neuropilin family, is a high affinity receptor for the semaphorins Sema E and Sema IV but not Sema III." *Neuron* 19, 547-559). However, the short cytoplasmic tail of neuropilins seems to be dispensable for their biological activity (Nakamura, et al (1998) *supra*), indicating the requirement of an associated co-receptor for signal transduction. Interestingly, in *Drosophila* (where neuropilins have not been identified to date) Plexin A is sufficient to mediate the biological response to semaphorin-1 in axon guidance (Winberg et al., 1998 *supra*).

In an initial set of experiments, we could not observe binding of the class 3 semaphorins Sema3A(Sema III), Sema3C(Sema E) or Sema3F(Sema IV) to plexins-A1, A2, A3, B1, B2 or C1 (not shown). To test whether plexins might be coreceptors with neuropilins for class 3 semaphorins, we set up co-precipitation experiments in COS cells to test whether neuropilins may interact with plexins. Three tested plexins (plexin-A1, -A3 and -B1) associated both with neuropilin-2 (Np2, shown in Fig. 3) and neuropilin-1 (not shown). The binding was specific, inasmuch as neither neuropilin nor any plexins coimmunoprecipitated with the netrin receptor DCC, under conditions

where DCC coimmunoprecipitated with the other netrin receptor UNC5H2 (Fig. 3 and data not shown). We observed finally that the plexin-neuropilin association is mediated by the *sema domain* of plexins, as demonstrated using either the “plexin-B1 truncated” splice variant (Figure 3) or an even shorter form of the extracellular domain (“plexin-B1-sema”, see Experimental procedures, not shown).

To further support the idea of a plexin-neuropilin multimeric receptor complex for semaphorins, we show here that plexin-A3 (e.g.) is expressed in a large number of neuronal classes, including sensory, sympathetic, motor, and olfactory bulb neurons (Figure 4 and data not shown), which are known to respond to class 3 semaphorins, and which express either neuropilin-1 or neuropilin-2 or both (Chen et al., 1997 *supra*; Feiner, L., Koppel, A.M., Kobayashi, H., and Raper, J.A. (1997). Secreted chick semaphorins bind recombinant neuropilin with similar affinities but bind different subsets of neurons in situ. *Neuron* 19, 539-545; He and Tessier-Lavigne, 1997 *supra*; Kolodkin et al., 1997 *supra*). Thus, plexin-A3 is a candidate for a physiological coreceptor involved in mediating class 3 semaphorin effects on these axons. Other plexins may also have a role as neuropilin coreceptors in specific cell populations, such as plexin-A2, which is expressed in a subset of sensory neurons and in dorsal horn cells, and plexin-A1, which is expressed at low levels and broadly in the spinal cord (Figure 4).

To directly test the possible involvement of plexins in class 3 semaphorin signal transduction, we studied the repulsive responses of *Xenopus* spinal neurons to *Sema3A*, which is mediated by a receptor mechanism involving neuropilin-1 (Song et al., 1998 *supra*). We asked whether these responses could be altered by expression of a presumed dominant-negative plexin-A1 construct lacking the cytoplasmic domain of the protein. Transmembrane proteins can be reliably expressed in these neurons by injecting the encoding mRNA at the developmental two cell stage, allowing the embryos to grow to tadpole stage, and then removing the spinal cord and culturing the neurons (Hong et al., 1999 *supra*). We therefore injected the mRNA encoding the truncated plexin-A1 construct, together with mRNA encoding GFP (as a reporter) and then studied the responses of spinal neurons expressing GFP that were derived from these embryos. Whereas control spinal neurons are repelled by *Sema3A* (Figure 5A, B and Song et al. 1998 *supra*), neurons from embryos injected with mRNA for truncated plexin-A1 did not respond with either repulsion or attraction to *Sema3A* (Figure 5C,

D). This blocking effect appeared to be specific, since expression of a different heterologous receptor, UNC5H2, did not impair repulsion by Sema3A (Hong et al., 1997 *supra*), and since expression of the truncated plexin construct did not block attractive responses to netrin-1 (Figure 5E, F). Figure 5G, H quantifies these effects.

- 5 As can be seen, the effect of Sema3A is completely abolished by the truncated plexin; although there is a slight apparent decrease in the attractive effect of netrin-1 the effect is not statistically significant.

Although we have used a truncated plexin-A1 construct, this construct may be expected to interfere with the function of various plexins, since all the plexins tested
10 (A1, A3 and B1) associated with neuropilin-1. These results support a role for one or more plexins in mediating the repulsive Sema3A signal in the *Xenopus* spinal neurons.

Example 10

Plexins signal via a novel type of tyrosine phosphorylated cytoplasmic domain

The sequences of plexin cytoplasmic domains are highly conserved among
15 plexins but do not match any known sequences. We found that the plexin-A3 and plexin-B1 proteins are phosphorylated on tyrosine residues when overexpressed in human kidney cells (BOSC-23), as demonstrated using anti-phosphotyrosine antibodies (Fig. 6a). Furthermore, after immunoprecipitation and in vitro kinase assays, plexin-A3 and plexin-B1 became phosphorylated (Fig. 6b). Resistance to an alkali treatment (see
20 Experimental procedures) confirmed the specific phosphorylation of tyrosine residues.

The cytoplasmic domains of several receptors, including Met proteins, become tyrosine phosphorylated owing to an intrinsic kinase activity (Ullrich, A. and Schlessinger, J. (1990) "Signal transduction by receptors with tyrosine kinase activity." *Cell* 61, 203-212). Since the cytoplasmic domain of plexins is not similar to any bona
25 fide or atypical tyrosine kinase, this suggests that a distinct tyrosine kinase co-immunoprecipitates in association with plexins, and is responsible for their tyrosine phosphorylation. Although some additional phosphorylated proteins can be found specifically with plexin-A3 and -B1, we have not as yet identified this associated kinase. A number of endogenously expressed tyrosine kinases, namely Met, Ron, Abl
30 and Src, were not found associated with plexin-A3 by immunoprecipitation and Western blotting (not shown). Since tyrosine phosphorylated residues often function as docking sites for intracellular signal transducers (Cantley, L.C., Auger, K.R., Carpenter, C., Duckworth, B., Graziani, A., Kapeller, R., and Soltoff, S. (1991) "Oncogenes and

signal transduction." Cell 64, 281-302), the fact that the cytoplasmic domains of plexins are tyrosine phosphorylated further suggests that they are part of signaling complexes.

Example 11

5 Plexin-A3 expressing cells induce repulsion of co-cultured cells

Stable transfectants expressing recombinant human plexin-A3 were successfully obtained in four different cell lines: IMR32 and AF8 (human neuroblasts), and BOSC-23 and MDCK (human and canine kidney cells, respectively). We observed modest phenotypic changes in the transfected cells, which generally become flatter and larger in size. The growth rate of plexin-A3 overexpressing cells was comparable to parental lines and we did not observe differences in the ability to adhere on different substrates (data not shown).

In keeping with previous report on the related Plexin of *Xenopus laevis* (Ohta, K., Mizutani, A., Kawakami, A., Murakami, Y., Kasuya, Y., Takagi, S., Tanaka, H., and Fujisawa, H. (1995). "Plexin: a novel neuronal cell surface molecule that mediates cell adhesion via a homophilic binding mechanism in the presence of calcium ions." Neuron 14, 1189-1199), we observed a modest increase in calcium-dependent homotypic cell aggregation of plexin-A3 transfectants (not shown). Surprisingly, we found that epithelial MDCK cells overexpressing plexin-A3 mediate strong repelling cues for adjacent cells. This was observed by co-culturing mock-transfected and plexin-A3 overexpressing MDCK cells together with several non-epithelial cell lines (such as NIH3T3, KJ29, and D17; Fig. 7A). Mock MDCKs grew alongside mesenchymal cells until confluency, when both cell types stopped proliferating. In contrast, when plexin-A3-overexpressing epithelial cells were grown in the same conditions, the adjacent mesenchymal cells withdrew from them, and ultimately detached from the plate.

To analyze the dynamics of this repulsion process, we monitored for 36 hours, by time-lapse video-microscopy, mixed cultures of transfected MDCK cells and fibroblasts, in a number of independent experiments. At low cell density, fibroblasts showed intrinsic motility, exploring the surface of the plate with long lamellipodia and filopodia, and thus coming in contacts with a high number of stationary MDCK islets. The time-length of the contacts between fibroblasts and control MDCK cells varied from 30 minutes to several hours, lasting mostly over 100 minutes. However, when fibroblasts were cultured with MDCK cells overexpressing plexin-A3, transient

contacts were observed, often lasting less than 30 minutes (see Fig. 7C). At higher cell density, fibroblasts stopped and clustered alongside the islands of control MDCKs, whereas they kept moving in a hectic fashion between the islands of plexin-A3 transfected cells (data not shown).

5 This cell-repelling effect is not due to the release of soluble factors, since exchanging conditioned media between mixed cultures was without effect (not shown). Moreover, the two different cell populations grew normally until they came into contact, indicating that the repelling effect requires cell-cell interaction. To rule out the possibility that plexin-A3 expressing cells generate an apoptotic signal for fibroblasts,
10 we monitored cell viability and apoptosis by TUNEL staining. As shown in Figure 7B, the clusters of repelled fibroblasts did not include apoptotic cells; furthermore, the detaching cells still excluded Trypan blue stain and were able to spread again on a new culture plate (not shown).





Taken together, these results demonstrate that in our experimental system, plexin-
15 A3 mediates cell repelling cues, presumably by interacting with surface bound ligands on opposing cells. We could not identify –so far– the specific ligand for plexin-A3, however we propose that this may be a transmembrane semaphorin. It should be noted that the intracellular domains of transmembrane semaphorins, such as Sema4D, also include tyrosine residues, which may themselves become phosphorylated and associate
20 with cytoplasmic signal transducer molecules, a property shown for ligands of the ephrin family (Holland, S.J., Gale, N.W., Mbamalu, G., Yancopoulos, G.D., Henkemeyer, M., and Pawson, T. (1996) "Bidirectional signalling through the EPH-family receptor Nuk and its transmembrane ligands." *Nature* 383, 722-725).

What is claimed is:

- 5 1. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence that encodes an amino acid sequence selected from the group consisting of the amino acid sequence shown in (SEQ ID NO: 2 (plexin B-2)), (SEQ ID NO: 4 (plexin B-3)), (SEQ ID NO: 6 (plexin D-1)) and (SEQ ID NO: 8 (plexin A-4))
- 10 2. Isolated nucleic acid having at least 80% nucleic acid sequence identity to a nucleotide sequence selected from the group consisting of the nucleotide sequence shown (SEQ ID NO: 1 (plexin B-2)), (SEQ ID NO: 3 (plexin B-3)), (SEQ ID NO: 5 (plexin D-1)) and (SEQ ID NO: 7 (plexin A-4)).
3. A vector comprising the nucleic acid of any one of claims 1 or 2.
- 15 4. An isolated polypeptide having at least 80% amino acid sequence identity to an amino acid sequence selected from the group consisting of the amino acid sequence shown in (SEQ ID NO: 2 (plexin B-2)), (SEQ ID NO: 4 (plexin B-3)), (SEQ ID NO: 6 (plexin D-1)) and (SEQ ID NO: 8 (plexin A-4)).
5. An isolated polypeptide having at least 80% amino acid sequence identity to:
 - 20 (a) the polypeptide shown in (SEQ ID NO: 2 (plexin B-2)), (SEQ ID NO: 4 (plexin B-3)), (SEQ ID NO: 6 (plexin D-1)) and (SEQ ID NO: 8 (plexin A-4)), lacking its associated signal peptide;
 - (b) an extracellular domain of the polypeptide shown in (SEQ ID NO: 2 (plexin B-2)), (SEQ ID NO: 4 (plexin B-3)), (SEQ ID NO: 6 (plexin D-1)) and (SEQ ID NO: 8 (plexin A-4)), with its associated signal peptide; or
 - 25 (c) an extracellular domain of the polypeptide shown in (SEQ ID NO: 2 (plexin B-2)), (SEQ ID NO: 4 (plexin B-3)), (SEQ ID NO: 6 (plexin D-1)) and (SEQ ID NO: 8 (plexin A-4)), lacking its associated signal peptide.

6. A chimeric molecule comprising a polypeptide according to claim 4 or 5 fused to a heterologous amino acid sequence.
7. The chimeric molecule of claim 6, wherein the heterologous amino acid sequence is a Fc region of an immunoglobulin.
- 5 8. An antibody that specifically binds to a polypeptide according to claim 4 or 5.
9. The antibody according to claim 8, wherein the antibody is a monoclonal, a humanized antibody or a single-chain antibody.
- 10 10. A method of suppressing or altering aberrant cell growth involving a signaling pathway between a plexin and a neuropilin in a mammal comprising the step of administering an effective amount of an agent to said mammal capable of interfering with the association between the plexin and neuropilin
- 15 11. A method of treating, suppressing or altering a disorder involving aberrant immune regulation involving a signaling pathway between a plexin and a neuropilin in a mammal comprising the step of administering an effective amount of an agent to said mammal capable of interfering with the association between the plexin and neuropilin.
- 20 12. The method according to claim 10 or 11 wherein said agent is a chimeric molecule according to claim 6 or 7.
13. The method according to claim 10 or 11, wherein said agent is an antibody according to claim 8 or 9.
- 25 14. A method of diagnosing or screening for tumors in a subject characterized by the expression profiles of the polypeptides according to claim 4 or 5 wherein the expression profile of the polypeptides is different in a non-tumor sample as compared to the expression profile of the polypeptides in a tumor sample.

2/7



F162

3/7




FIG. 3

4/7




Fig. 4

5/7









FIG. 5



7/7
FIG. 7

SEQUENCE LISTING

<110> University of Torino

<120> Novel Plexins and Uses Thereof

<130> A077PCT

<140> Not assigned yet

<141> 2000-08-25

<150> 60/150576

<151> 1999-08-25

<160> 12

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 6252

<212> DNA

<213> HOMO SAPIEN

<400> 1

gcggggggca	atggcaactgc	agctctgggc	cctgaccctg	ctgggacctg	tgggcgcagg	60
tgccagcctg	aggccccgca	agctggactt	cttcgcgagc	gagaaaagagc	tgaaccacct	120
ggctgtggat	gaggcctcag	gcgtgggtgta	cctggggggc	gtgaatgcc	tctaccagct	180
ggatgcgaag	ctgcagctgg	agcagcaggt	ggccacgggc	ccggccctgg	acaacaagaa	240
gtgcacgcgc	cccatcgagg	ccagccagtg	ccatgaggct	gagatgactg	acaatttcaa	300
ccagctgtct	ctgctcgacc	ctcccgagaa	gcgcttggtg	gagtgccgca	ccctcttcaa	360
gggcctctgc	gctctgcgcg	ccctgagcaa	catctccctc	cgctgttctt	acgaggacgg	420
cagcggggag	aagtcttctg	tgcccgagaa	tgatgagggc	gtggccacag	tggggctggt	480
gagctccacg	ggtcctgggt	gtgaccgcgt	gctgttttgt	ggcaaaagca	atggggccaca	540
cgacaacgcg	atcatcgatg	gcactcggtt	gttggaccgg	actgacagca	gggaggccctt	600
tgaagcctac	acggaccacg	ccacctacaa	ggccggctac	ctgtccacca	acacacagca	660
gttcgtggcg	gccttcgagg	acggccccta	cgtcttcttt	gtcttcaacc	agcaggacaa	720
gcaccgggcc	cggaaccgca	cgctgctggc	acgcatgtgc	agagaagacc	ccaactacta	780
ctctaccctg	gagatggacc	tgcatgtccg	ggaccccgac	atccacgccg	ctgcctttgg	840
cacctgcctg	gcgcgctccg	tggtgctggc	tggtctctgg	agggtgctat	atgctgtctt	900
cagcagagac	agccggagca	gtggggggcc	cggtgcgggc	ctctgcctgt	tcccgctgga	960
caaggtgcac	gccaaagatg	aggccaaccg	caacgcctgt	tacacagca	cccgggaggc	1020
ccgtgacatc	tctacaagc	ccttccacgg	cgatatccag	tgccggcgcc	acgcgcgggg	1080
ctccagcaag	agcttcccat	gtggctcgga	gcacctgcc	taccgcctgg	gcagcccgga	1140
cgggctcaga	ggcacagcgc	tgtctgcagc	tggaggccctg	aaacctcagc	cgctgacggt	1200
gcgcgcgcag	aacaaccaca	ctgttgcctt	tctgggcacc	tctgatggcc	ggatccctcaa	1260
ggtgtacatc	acccccagatg	gcacctctct	agagtacgac	tctatccttg	tgagataaaa	1320
caagagagtc	aagcgcgacc	tggtactgtc	tggagacctg	ggcagccctg	acgccatgac	1380
ccagagacag	gtgttctccg	tgccggtgca	ggagtgcctg	agctacccca	cctgcaccca	1440
gtgcccgcgac	tcccaggacc	cctactgcgg	ctgggtgcgt	gtcgaggagc	gatgcacccc	1500
gaagggcagc	tgtccgcggg	ccgaggaggc	cagccaactgg	ctgtggagcc	gaagcaagtc	1560
ctgcgtggcg	gtcaccagcg	cccagccaca	gaacatgagc	cggcgggccc	agggggaggt	1620
gcagctgacc	gtcagccccc	tccctgcctt	gagcgaggag	cagcagttgc	tgtgcctttt	1680
tggggagtgc	ccgccacacc	ccgcccgcgt	ggagggcgag	gccgtcatct	gcaactcccc	1740

aagcagcatt	cccgctacac	gcggcaggcca	ggaccacgtg	gccgtgacca	tcacagctcct	1800
ccttagacga	ggcaacatct	tcctcacgtc	ctaccagttac	cccttctacg	actgcccgca	1860
ggccatgagc	gtggaggaga	acctgcgctg	catctccctgc	gtgagcaacc	gctggacctg	1920
ccagctggagc	ctgcgctacc	acaggtgcgc	ggaggctctg	cccaaccctg	aggagcgcat	1980
cgtccgtggc	cacatggagg	acagctgtcc	ccagttccctg	ggaccaccgc	cctcgggtgat	2040
ccccatgaac	cacgagacag	atgtgaacct	ccaggggcaag	aaacctggaca	ccgtgaaggg	2100
ttcctccctg	cagctgggca	gtgacttgct	caagttcatg	gagccggtga	ccatgcaggga	2160
atccgtggacc	ttcgctcttc	ggaccccaaaa	gctgtccacc	gatgcccaag	agacgtgtgc	2220
cccgccactc	tacgtcaagt	cttacggcaa	gaatatcgac	agcaagctcc	atgtgacctt	2280
ctacaactgc	ctctcttgcc	gcagcgactg	cagcctgtgc	cgggcccgta	accgcgacta	2340
cagggtggcg	tggtgcgggg	gccagagcag	gtgcgtgtat	gagagccctg	gcaacaccac	2400
ctccagctgc	ccgcgcggcg	tcattcaacag	gatccagcct	gagacggggc	ccctgggtgg	2460
gggcaaccgc	atcaacatcc	tggggtccaa	tttgggctgc	caagcagggg	acatccagag	2520
gcctctgtg	ccgcgcggga	actgctcttt	tcagccggaa	cgttactccg	ttgtccaccg	2580
gatccgtgtg	gtgatcgagg	ctcgggagac	gccttccacg	gggggtgtcg	agggtgagct	2640
ctccgggaac	ctgggcgctt	cgctcccaa	tgtccagttc	accttccaac	agcccaagcc	2700
ttccatgtg	gagccgcacg	agggaccgca	ggcggcgccg	accacactga	ccatccagcc	2760
caccacactg	gcacacggct	ccagagagga	cgctcggtgt	accctcaacg	gcgtcccggt	2820
taaaagtacc	aagtctgggg	cgacgtccca	gtgtgtcact	ggccccagag	gcacacgggg	2880
ccagatgct	ctggaggtct	ctacgggggg	gtccccctgt	cccaaccctg	gcattctctt	2940
ccactaccgc	gaaaaaccgc	tactgcgagc	cttcgagccg	ctacgaagct	ttgcagctgt	3000
tgcccgctgc	atacaactga	cggtgcaggg	cttcagcctg	atccagaggt	ttggcatggt	3060
ggctcatcgc	gagccccctg	agtcctggca	ccgcgcggcg	gaggttgaa	ttgcagctgc	3120
cattcagctg	gtgggtacag	atacgtgtt	ccacaatgac	accaaggtct	ctcttcctgc	3180
cccgctctgt	ctctgaggagc	cagaggcccta	caacctcacg	gtgctgatgc	agatggagcg	3240
gcacctctgc	ctgctcagaa	cagaggccgg	ggccttcgag	tacgtgtcag	acccccactt	3300
tgagaacttc	acaggtggcg	tcaagaagca	ggtcaacaag	ctcatccacg	ccccggggc	3360
caatctgaac	aagggcattg	cgtctcagga	ggccgaggcc	ttcgtgggtg	ctcccgactg	3420
caccatgaag	acgtctgacg	agaccgacct	gtactgtgag	cccccgagg	tgacgcccc	3480
gccccagcgc	cgccagaaag	gagacaccac	acacaacctg	cccgagttca	ttgtgaagtt	3540
cggtctctgc	gagtggtgtc	tgggcccgct	ggagtacgac	acacgggtga	gcgacgtgct	3600
gctcagcttc	atcttgcgcg	tggtcatcgt	gccatgtgtg	gtcgtctagt	cggtgtggtg	3660
ctactgtctac	tgaggagaaga	gccacgaggg	cgaaacgagag	tatgagaaga	tcaagtccca	3720
gctggaggagc	ctggaggaga	cggtgcggga	ccgctgcaag	aagggaattca	cagacctgat	3780
ctcgagattac	gaggaccaga	ccaacgacgt	gcacgaggcc	ggcatccccg	tgctggacta	3840
caagacctac	accgacggc	ttcttctctc	gccctccaag	gacggcgaca	aggacgtgat	3900
gcacaccggc	aagctggaca	tccttgagcc	cgccggggcg	gtggtggagc	aggccctcta	3960
ccagttctcc	aaactctgcg	acagcaagtc	tttctctatc	aatttctatc	acaccttgga	4020
gaaccagcgg	gagttctcgg	ccccgcgcaa	ggtctacttc	gcgtcccttc	tgacgtggcg	4080
gctgcacggg	aaactggagt	actacacgga	catcatgcac	acgctctctc	tgagactctc	4140
ggagcagttac	gtggtggcca	agaaccccaa	gctgatgctg	cgcaggtctg	agactgtggt	4200
ggagaggtat	ctgtccaaat	ggatgtccat	ctgcctgtac	cagtaacctc	gacacagttg	4260
cggggagccc	ctgtcaaacg	tcttcaaggg	catcaaacat	caggtggaaa	agggcccggt	4320
ggatgcggtg	cagaagaagg	ccaagtacac	tctcaacgac	acggggctcg	tgggggatga	4380
tgtagagtac	gcacccctga	cggtgagcgt	gatcgtgcag	gacgagggag	tggaacgcat	4440
ccccgtgaag	gtctctaac	gtgacacatt	ctccaggttc	aaggagaaga	ctattgacca	4500
gggtacccgt	ggggcagcct	gtctctctgt	gccacggcca	gacagcgtgg	tccttgaggtg	4560
gcgtgtccag	tcacacggc	agatccgttc	ggactctggc	ctgacgtcac	agcgggaggg	4620
ccggttggaag	cgctgaacca	cccttatgca	ctacaattgt	cggtatggag	ccacctctat	4680
ccgtgtccaa	ttgggggtct	ccacgagcc	ggaggacagc	cagcagacc	tgcttgggga	4740
gggccatgac	ctcttgaggg	aggagaaccg	ggtgtggcac	ctggtgcggc	cgaccgcagca	4800
ggcgaagtc	ggcaagttca	agagagggc	cgtaaaagag	aaggagcggg	cgaaagccat	4860
caccagatgc	tacctgacgc	ggctgccttc	agtcgaaggg	acactgcagc	agttttggga	4920
caactctctc	cagagcgtgc	tgccgcctgt	gcacgcgggt	ccacctgcag	tcaagtaatt	4980
cttcgacttc	ctggcagagc	agggcagagaa	gcacaacatc	caggatgaag	acaccattcca	5040

catctggaag	acgaacacgt	taccgctccg	gtcttgggtg	aacatctcta	agaaccccc	5100
cttcattctt	gacgtgcatg	tccacgaggt	gggtggacgcc	tcgctgtcag	tcacgcgcga	5160
gaccttcttg	gatgcctgca	cgcgcacgga	gcataagctg	agccgcgatt	ctcccagcaa	5220
caagctgctg	tacgccaagg	agatctccac	ctacaagaag	atggtggagg	ataactacaa	5280
ggggatcctg	cagatgggtc	aggctcagca	ccaggacatg	aacacacacc	tggcagagat	5340
ttcccgggcg	cacacggact	ccttgaacac	cctcgtggca	ctccaccagc	tctaccaata	5400
cacgcagaag	tactatgacg	agatcatcaa	tgccctggag	gaggaacctg	cgccccagaa	5460
gatgcagctg	gccttcgccg	tgacgcagat	tgccgctgca	ctggagaaca	aggctactga	5520
cctctgacct	acaatctcca	gtgctgcctt	gggacatagg	tacctgaggt	acctgagagc	5580
ccctcagggg	aggaggcgca	gtggctgtgg	ctgaggcccc	cacctctccc	tggaaacgcg	5640
cccaagccgg	agtgggtgca	gcccgaaccc	gcccagcgtc	tagactgtag	catcttcttc	5700
tgagcaatac	cgccggggcac	cgcaccagca	ccagcccccag	ccccagctcc	ctccggccgc	5760
agaaccagca	tcgggtgttc	actgtcagat	ctcagatgat	tgtaaaatgt	cgcttacgct	5820
gccacgctgg	gggcagctgg	cctccgcctc	cgcccacgca	ccagcagccg	cctccatgcc	5880
ctaggttggg	ccctcggggg	atctgagggc	ctgtggcccc	caggggcaagt	tcccagatcc	5940
tatgtctgtc	tgctcaccac	gagatggggg	gaggagaaaa	agcggtagca	tgcccttctg	6000
acctcaccgg	ccctccccaag	gggtcccgca	ctctgggtgg	actcacggct	gctgggcccc	6060
acgtcaaaag	tcaagttaga	cgtaggtcaa	gtctctacgt	ggggcccaga	catcttgggg	6120
tcctgtgtctg	tcagacagcc	tgccctagag	ccccaccag	ctcgggggga	ctggggagag	6180
ttccaagacc	accaccaacc	tttttgtaaa	tcttgttcat	tgtaaatcaa	atacagcgct	6240
tttttactc	cg					6252

<210> 2

<211> 1838

<212> PRT

<213> HOMO SAPIEN

<400> 2

Met	Ala	Leu	Gln	Leu	Trp	Ala	Leu	Thr	Leu	Leu	Gly	Leu	Leu	Gly	Ala
1				5					10					15	
Gly	Ala	Ser	Leu	Arg	Pro	Arg	Lys	Leu	Asp	Phe	Phe	Arg	Ser	Glu	Lys
			20					25					30		
Glu	Leu	Asn	His	Leu	Ala	Val	Asp	Glu	Ala	Ser	Gly	Val	Val	Tyr	Leu
		35					40				45				
Gly	Ala	Val	Asn	Ala	Leu	Tyr	Gln	Leu	Asp	Ala	Lys	Leu	Gln	Leu	Glu
	50					55					60				
Gln	Gln	Val	Ala	Thr	Gly	Pro	Ala	Leu	Asp	Asn	Lys	Lys	Cys	Thr	Pro
65				70					75					80	
Pro	Ile	Glu	Ala	Ser	Gln	Cys	His	Glu	Ala	Glu	Met	Thr	Asp	Asn	Val
			85					90					95		
Asn	Gln	Leu	Leu	Leu	Leu	Asp	Pro	Pro	Arg	Lys	Arg	Leu	Val	Glu	Cys
			100				105						110		
Gly	Ser	Leu	Phe	Lys	Gly	Ile	Cys	Ala	Leu	Arg	Ala	Leu	Ser	Asn	Ile
		115				120					125				
Ser	Leu	Arg	Leu	Phe	Tyr	Glu	Asp	Gly	Ser	Gly	Glu	Lys	Ser	Phe	Val
	130					135					140				
Ala	Ser	Asn	Asp	Glu	Gly	Val	Ala	Thr	Val	Gly	Leu	Val	Ser	Ser	Thr
145				150				155						160	
Gly	Pro	Gly	Gly	Asp	Arg	Val	Leu	Phe	Val	Gly	Lys	Gly	Asn	Gly	Pro
			165					170					175		
His	Asp	Asn	Gly	Ile	Ile	Val	Ser	Thr	Arg	Leu	Leu	Asp	Arg	Thr	Asp
		180					185					190			
Ser	Arg	Glu	Ala	Phe	Glu	Ala	Tyr	Thr	Asp	His	Ala	Thr	Tyr	Lys	Ala

Gly	Tyr	195				200				205					
		Leu	Ser	Thr	Asn	Thr	Gln	Gln	Phe	Val	Ala	Ala	Phe	Glu	Asp
Gly	Pro	Tyr	Val	Phe	Phe	Val	Phe	Asn	Gln	Gln	Asp	Lys	His	Pro	Ala
225	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Tyr	Ser	Ala	Ala	Phe	Gly	Thr	Cys	Leu	Ala	Arg	Ser	Val	Ala	Pro	Gly
250	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Tyr	Ser	Ala	Ala	Phe	Gly	Thr	Cys	Leu	Ala	Arg	Ser	Val	Ala	Pro	Gly
275	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Tyr	Ser	Ala	Ala	Phe	Gly	Thr	Cys	Leu	Ala	Arg	Ser	Val	Ala	Pro	Gly
290	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Gly	Gly	Pro	Gly	Ala	Gly	Leu	Cys	Leu	Phe	Pro	Leu	Asp	Lys	Val	His
305	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Ala	Lys	Met	Glu	Ala	Asn	Arg	Asn	Ala	Cys	Thr	Thr	Gly	Thr	Arg	Glu
320	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Gly	His	Ala	Pro	Gly	Ser	Ser	Lys	Ser	Phe	Pro	Cys	Gly	Ser	Glu	His
335	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Leu	Pro	Tyr	Pro	Leu	Gly	Ser	Lys	Ser	Phe	Pro	Cys	Gly	Ser	Glu	His
350	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Leu	Gln	Arg	Gly	Gly	Leu	Asn	Leu	Thr	Ala	Val	Thr	Val	Ala	Ala	Glu
365	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Leu	Pro	Tyr	Pro	Leu	Gly	Ser	Lys	Ser	Phe	Pro	Cys	Gly	Ser	Glu	His
380	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Leu	Gln	Arg	Gly	Gly	Leu	Asn	Leu	Thr	Ala	Val	Thr	Val	Ala	Ala	Glu
395	Arg	Asn	Arg	Thr	Leu	Ala	Arg	Met	Cys	Arg	Glu	Asp	Pro	Asn	Thr
Lys	Val	Tyr	Leu	Thr	Pro	Asp	Gly	Thr	Ser	Ser	Glu	Tyr	Asp	Ser	Ile
410	Arg	Asn	Arg	Thr											

Leu Arg Tyr His Glu Cys Arg Glu Ala Ser Pro Asn Pro Glu Asp Gly
 645
 Ile Val Arg Ala His Met Glu Asp Ser Cys Pro Gln Phe Leu Gly Pro
 660
 Ser Pro Leu Val Ile Pro Met Asn His Glu Thr Asp Val Asn Phe Gln
 675
 Gly Lys Asn Leu Asp Thr Val Lys Gly Ser Ser Leu His Val Gly Ser
 690
 Asp Leu Leu Lys Phe Met Glu Pro Val Thr Met Gln Glu Ser Gly Thr
 705
 Phe Ala Phe Arg Thr Pro Lys Leu Ser His Asp Ala Asn Glu Thr Leu
 720
 Pro Leu His Leu Tyr Val Lys Ser Tyr Gly Lys Asn Ile Asp Ser Lys
 740
 Leu His Val Thr Leu Tyr Asn Cys Ser Phe Gly Arg Ser Asp Cys Ser
 755
 Leu Cys Arg Ala Ala Asn Pro Asp Tyr Arg Cys Ala Trp Cys Gly Gly
 770
 Gln Ser Arg Cys Val Tyr Glu Ala Leu Cys Asn Thr Thr Ser Glu Cys
 785
 Pro Pro Pro Val Ile Thr Arg Ile Gln Pro Glu Thr Gly Pro Leu Gly
 800
 Gly Gly Ile Arg Ile Thr Ile Leu Gly Ser Asn Leu Gly Val Gln Ala
 820
 Gly Asp Ile Gln Arg Ile Ser Val Ala Gly Arg Asn Cys Ser Phe Gln
 835
 Pro Glu Arg Tyr Ser Val Ser Thr Arg Ile Val Cys Val Ile Glu Ala
 850
 Ala Glu Thr Pro Phe Thr Gly Gly Val Glu Val Asp Val Phe Gly Lys
 865
 Leu Gly Arg Ser Pro Pro Asn Val Gln Phe Thr Phe Gln Gln Pro Lys
 885
 Pro Leu Ser Val Glu Pro Gln Gln Gly Pro Gln Ala Gly Gly Thr Thr
 900
 Leu Thr Ile His Gly Thr His Leu Asp Thr Gly Ser Gln Glu Asp Val
 915
 Arg Val Thr Leu Asn Gly Val Pro Cys Lys Val Thr Lys Phe Gly Ala
 930
 Gln Leu Gln Cys Val Thr Gly Pro Gln Ala Thr Arg Gly Gln Met Leu
 945
 Leu Glu Val Ser Tyr Gly Gly Ser Pro Val Pro Asn Pro Gly Ile Phe
 965
 Phe Thr Tyr Arg Glu Asn Pro Val Leu Arg Ala Phe Glu Pro Leu Arg
 980
 Ser Phe Ala Ser Gly Gly Arg Ser Ile Asn Val Thr Gly Gln Gly Phe
 995
 Ser Leu Ile Gln Arg Phe Ala Met Val Val Ile Ala Glu Pro Leu Gln
 1010
 Ser Trp Gln Pro Pro Arg Glu Ala Glu Ser Leu Gln Pro Met Thr Val
 1025
 Val Gly Thr Asp Tyr Val Phe His Asn Asp Thr Lys Val Val Phe Leu
 1045
 Ser Pro Ala Val Pro Glu Glu Pro Glu Ala Tyr Asn Leu Thr Val Leu
 1060
 Ile Glu Met Asp Gly His Arg Ala Leu Leu Arg Thr Glu Ala Gly Ala
 1070
 1085

[illegible]

Ser Thr Ala Gln Ile Leu Ser Asp Leu Asp Leu Thr Ser Gln Arg Glu
 1525 1530 1535
 Gly Arg Trp Lys Arg Val Asn Thr Leu Met His Tyr Asn Val Arg Asp
 1540 1545
 Gly Ala Thr Leu Ile Leu Ser Lys Val Gly Val Ser Gln Gln Pro Glu
 1555 1560 1565
 Asp Ser Gln Gln Asp Leu Pro Gly Glu Arg His Ala Leu Leu Glu Glu
 1570 1575 1580
 Glu Asn Arg Val Trp His Leu Val Arg Pro Thr Asp Glu Val Asp Glu
 1585 1590 1595 1600
 Gly Lys Ser Lys Arg Gly Ser Val Lys Glu Lys Glu Arg Thr Lys Ala
 1605 1610 1615
 Ile Thr Glu Ile Tyr Leu Thr Arg Leu Leu Ser Val Lys Gly Thr Leu
 1620 1625 1630
 Gln Gln Phe Val Asp Asn Phe Phe Gln Ser Val Leu Ala Pro Gly His
 1635 1640 1645
 Ala Val Pro Pro Ala Val Lys Tyr Phe Phe Asp Phe Leu Asp Glu Gln
 1650 1655 1660
 Ala Glu Lys His Asn Ile Gln Asp Glu Asp Thr Ile His Ile Trp Lys
 1665 1670 1675 1680
 Thr Asn Ser Leu Pro Leu Arg Phe Trp Val Asn Ile Leu Lys Asn Pro
 1685 1690 1695
 His Phe Ile Phe Asp Val His Val His Glu Val Val Asp Ala Ser Leu
 1700 1705 1710
 Ser Val Ile Ala Gln Thr Phe Met Asp Ala Cys Thr Arg Thr Glu His
 1715 1720 1725
 Lys Leu Ser Arg Asp Ser Pro Ser Asn Lys Leu Leu Tyr Ala Lys Glu
 1730 1735 1740
 Ile Ser Thr Tyr Lys Lys Met Val Glu Asp Tyr Tyr Lys Gly Ile Arg
 1745 1750 1755 1760
 Gln Met Val Gln Val Ser Asp Gln Asp Met Asn Thr His Leu Ala Glu
 1765 1770 1775
 Ile Ser Arg Ala His Thr Asp Ser Leu Asn Thr Leu Val Ala Leu His
 1780 1785 1790
 Gln Leu Tyr Gln Tyr Thr Gln Lys Tyr Tyr Asp Glu Ile Ile Asn Ala
 1795 1800 1805
 Leu Glu Glu Asp Pro Ala Ala Gln Lys Met Gln Leu Ala Phe Arg Leu
 1810 1815 1820
 Gln Gln Ile Ala Ala Leu Glu Asn Lys Val Thr Asp Leu
 1825 1830 1835

<210> 3

<211> 5367

<212> DNA

<213> HOMO SAPIEN

<400> 3

atggctctgc	ggcctccett	cggcctctgc	ctctctctgc	tgctgctgtc	cccaccgcca	60
ctggccttga	cagggggcca	tcgctctctc	gcaccttaata	ccactctcaa	ccacttgcca	120
ctggcaactg	gcgcaggaac	actctatgtc	ggcgcaagtga	accgcctctt	ccagctcagc	180
cccgagctgc	agctcgagagc	cgttgctgtc	actggccctg	taatcgacag	ccctgactgc	240
tgccctcttc	gtgaccagcc	cgagtgccca	caggccccagc	tcactgacaa	tgcccaaccag	300
ctgctgctgg	tgagcagccg	cgccagagg	ctggtggcct	cgggcgaggt	gcgcaggggc	360
gtgtgtgaga	cacggccctt	tggggatgtg	gccgaggtgc	tgtaccaggc	tgaggaccct	420
ggtgacgggc	agtttgtggc	tgccaatacc	ccgggagtg	caacgggtgg	gctggtggtg	480

cctctgccc	gcgaggac	cctgctgtg	gccagaggg	tgccgggcaa	gctgtcggca	540
gggggtgcc	ccttgccat	ccgcagctg	gccgggtctc	agccctcttc	cacgaggggc	600
ctgggcgcgc	tggtgtggg	cgactcttc	gaactacaac	acagctacgt	cggggctctt	660
gccgaagccc	gctccgccta	ctctgtgtc	cgccgcgcgc	gggcccgggc	ccaggctgag	720
taccctctct	acgtggccgc	cgctcgctg	gggggaacac	acctgtacac	ctacgtggag	780
gtccccctct	cctgcccagg	ccagggcctc	atccaggcgc	cctctcttgc	ccggggcacc	840
ttgctagggg	tgttttgcgc	gggcccgaag	ggcaccagag	cgctcaagtc	tgctctcccc	900
atggtggagg	tggtgtccag	catggagcag	gcccgaggac	ctgtctacac	ggcgggcggc	960
cgggggccca	gggggcgcag	ggaagccacc	gtggagtacg	gcgtcaagtc	ggcgtcgctc	1020
acccctgccc	ttgatctccc	cgagctgtac	cctctgtggc	acgagcacac	ccccagcccc	1080
attgctgggc	ggcagccctc	ggaggtccag	cctctgctga	agctcgggca	gccggtcagc	1140
gccgtggcag	ctctccaggc	agatgggcac	atgatagcct	tccctggggg	caccaggggc	1200
cagctgtaca	aggtcttctc	ccacggctcc	caggggccag	tttaccactc	caggcaagtg	1260
gggctccag	gctcagccat	cagcccagac	ctgtctgtgg	acagcagctg	cagtcacctc	1320
tatgtctga	ctgccaccac	ggtggaccgg	atacctgtgg	cagcctgcgc	ccaggtccct	1380
gaactgtcca	gctgctccca	ggcccaggac	ccgctgtgtg	gctgtgtgtg	cctccagggc	1440
agggtgtacc	ggaaggggca	gtg-gggcgg	cagggccagc	tgaaacagtg	gctgtggagt	1500
tatgaggagg	acagccactg	cttgacatc	cagagcctgc	tgccggggca	ccacccccgc	1560
caggagagca	ccagggtcac	tttgtctgtc	ccccgcctgc	ccatctgga	tgacagtga	1620
tacttccatt	gtgcgttcgg	ggactatgac	agccttggtc	atgtggaaag	gccccactgt	1680
cgtgtgtgca	ccccctccca	agaccagggt	ccaattaac	ctccagcgac	agaccactgc	1740
actgtccctg	tgccctgat	gttcaggagc	gtactgtgtg	ctgccaccaa	ctctctcttt	1800
tatgactgca	gtcccgctca	ggccttgagg	gcggctgcgc	ccgtctcttc	ccaggggcct	1860
ctgctcctct	tccactgtctg	gtggagcgtg	ctggagaac	ttcggggact	ccacccccac	1920
ctggaggagca	cagcagggga	ttcagccctc	atccactgcc	aggcccacac	cggggagctc	1980
ccagtgccca	tctacgtcac	ccagggtgaa	gccagaggcg	tgacacacac	ccatgtctct	2040
tatggtgagc	ctgaggggac	ccaggcaggc	ggggcagggt	gggtggcaga	caggaggcgc	2100
tcagacacat	gcctgaccct	cctagtgtat	ctgtacagac	tgcccatctg	gccaccggga	2160
ctgcagcaac	gtcccaagcg	ccaaacagg	cctgggtctc	ctgtgaccag	cctctgccca	2220
ggcccccaaa	ccccagcagc	tccgctctgg	tgggtctggt	ggctggccgg	gcaccacgca	2280
ctgcagatgt	gagcgtgggt	gcgggggacc	ccatctgcca	tcatttgctc	gctgcaggctc	2340
gagccccatg	ctgcctcccc	tgagggaagg	ttggccctca	ccatctctgg	ctccaaacctg	2400
ggcggccgct	tcgcgcatgt	gcagtaagcc	gaacctgtcc	tgtctgagct	gagctctcgc	2460
tgggggcccc	aggcaggggg	acaccagctc	accatccgag	gtcagcacct	ccagacaggt	2520
ggcaacacca	gtgccttcgt	gggtggccaa	cctgtccca	tggtggggcg	actgatccgt	2580
gacgcagagc	tgacagcttc	cagggccag	cgccctctac	tgctctgtgt	ctggagggtc	2640
ggagccctgt	ctgcgcaccc	ccagctctgt	atccagctcg	gtggggggct	gctgcagcgc	2700
acagcagagc	ccagctcact	ccactctgtg	tcggccctga	atgccccaca	gtgctccacc	2760
gtctgtctcg	tcaactctgtc	cagcctctct	ctgtgcggga	gcctgtctgt	accagacaga	2820
gccaccccgc	agcgggtctt	cttccacctc	gacacagctc	aagtggacct	cgccagtccc	2880
agtgggggca	agggtcttct	gtaccagccc	aacccccgcc	tgccacctct	cagccgcgag	2940
gggctctccc	gcccctaccg	cctcaagcca	ggccatgtcc	tggtgtgga	ggcgaggggc	3000
ctcaactctg	gcctacagcaa	ggaggagggt	cgctgcaca	tcggcccgcg	cgagtgcctg	3060
gtgaagagct	tcacgcgcac	ccactctgtc	tgccagccgc	ctgcgcagca	cccgagccct	3120
cctcaactctg	ccggcctgcc	acagttctgt	gtgcagatgg	gcaatgtcga	cgctggcctg	3180
ggccctgtgc	agtaagaggc	tgaacccccg	ctgtctgctc	ttccctgtgc	ggcccaggga	3240
ggcgtgggca	tggtgtctgc	agtgtgatt	gcggccgtgc	ctctctctac	cctcatgtac	3300
aggcacaaag	gcaagcagcg	cctgcgggac	taccagaagg	tgctagtcca	gctggagagc	3360
ctggagacac	gcgtgggaga	ccagtcggcg	aaggagtcca	cagacctcat	gacggagagt	3420
accagactca	gcagcgacct	ggaggggcagc	gggactccct	tcttggaacta	ccgcacctac	3480
gccagagctc	cctctctccc	tgcccatggc	ggttgcccg	tgaccgccaa	gcttgagggg	3540
ccaggggagg	acggccactg	tgccactgtg	cgccaggggc	tcacgcagct	ctccaaacctg	3600
ctcaacagca	agctcttctc	cctccagggt	agggccgtgt	cgccggagtg	ccagtgggc	3660
aaggagggtg	ggctggggaa	ctactggcct	gagacaaagg	tgggggaggga	gacagagacc	3720

atggttgaga	aactgtcac	caactggctg	tccactctgc	tgtacgcctt	cctgaggagg	3840
gtggctgggt	aaccactgta	catgctcttc	cgggccatcc	agtaccaggt	ggacaaaggc	3900
cccggtgacg	ccgtgacagg	caaggccaaa	cggaccctga	atgatagccg	cttgcctcgg	3960
gaggacgtgg	agttccagcc	cctgacgtg	atgggtgctg	tggggccggg	ggctggcggg	4020
gccgacggga	gcagcgagat	gcagcgcgtg	ccagcccggg	tgctcgacac	ggacaccatc	4080
acccaggtca	aggagaaggt	gttgaccac	gtctacaagg	gcaccccctt	ctcccagagg	4140
ccctcagtcg	atgccctaga	ccttggtag	agagccagcc	ctgccacccc	accccaggga	4200
ccctccctca	ccctccgggc	acctggagcc	cctcaactgt	gtcttactat	gaacataacc	4260
acgcctggag	atggcgagg	gggggggggt	tgctcttggc	acctggtgaa	agccaccgag	4320
gagccagaag	gggccaaggt	gcggtgcagc	agcctcgggg	agcgcgagcc	agcaagggcc	4380
aaggccattc	cggaatctta	cctcaccgt	ctgctgtcca	tgaaggttgg	tgcggcctgg	4440
gtggctgggg	ctgagaggag	gtcagccag	ggaccccgac	cgagccaggg	tgtgggaggg	4500
gcagggcgag	cctcagccgt	ggatggcccc	cacacccctg	cctccacaca	gcccttatcc	4560
cctgcctcgc	agggcagcgt	gcagaagttt	gtggacgaca	ccttccaggg	cattctcagc	4620
gtgaaccggc	ccatcccat	cgccgtcaag	tacctgtttg	accttctgga	tgagctagca	4680
gagaagctcg	gcacagagga	cccagggacc	ctgcacatct	ggaagaccaa	cagctctgctg	4740
ctcgggttct	gggtgaatgc	cttgaagaac	ccacagctca	cttttgatgt	acgggtgtcgc	4800
gacaatgtgg	acgccatctc	tgtctgtatc	gccacagact	tcattgactc	ctgtacacc	4860
tcgggacata	aagtggggcg	ggtgagagca	gtgccagcac	cagcagctgg	caggggcttg	4920
aggaggaag	gcttatgggg	gaagcctaga	gggctgtgca	cagagctctg	gggtggcagct	4980
ggcagcatca	tgggggcacc	ttcacctcgc	agctcatgcc	tagccctccc	ctctccctcg	5040
gagcaggatt	ccccagtgaa	caaacctgct	tacgcccggg	agatcccacg	ctacaaagcag	5100
atgggtggaga	ggatctatgc	ggacattcgc	cagagctctc	ggcgagatga	ccaggagatg	5160
aactctgctt	ggctgagct	ctccgggaac	tacacttctg	ctccccactg	tctggaggct	5220
ctgcaagaac	ctacaaacca	catccacagg	tactatgatc	agattatcag	ctcctggagg	5280
gaggaccctg	tggggcagaa	gctgcagctg	gcctgcgcgc	tgcagcaggt	cgccgccttg	5340
gtggaaaaca	aagtgactga	cctgtga				5367

<210> 4
 <211> 1788
 <212> FRT
 <213> HOMO SAPIEN

<400> 4
 Met Ala Arg Trp Pro Pro Phe Gly Leu Cys Leu Leu Leu Leu Leu Leu
 1 5 10 15
 Ser Pro Pro Pro Leu Pro Leu Thr Gly Ala His Arg Phe Ser Ala Pro
 20 25 30
 Asn Thr Thr Leu Asn His Leu Ala Leu Ala Pro Gly Arg Gly Thr Leu
 35 40 45
 Tyr Val Gly Ala Val Asn Arg Leu Phe Gln Leu Ser Pro Glu Leu Gln
 50 55 60
 Leu Glu Ala Val Ala Val Thr Gly Pro Val Ile Asp Ser Pro Asp Cys
 65 70 75 80
 Val Pro Phe Arg Asp Pro Ala Glu Cys Pro Gln Ala Gln Leu Thr Asp
 85 90 95
 Asn Ala Asn Gln Leu Leu Leu Val Ser Ser Arg Ala Gln Glu Leu Val
 100 105 110
 Ala Cys Gly Gln Val Arg Gln Gly Val Cys Glu Thr Arg Arg Leu Gly
 115 120 125
 Asp Val Ala Glu Val Leu Tyr Gln Ala Glu Asp Pro Gly Asp Gly Gln
 130 135 140
 Phe Val Ala Ala Asn Thr Pro Gly Val Ala Thr Val Gly Leu Val Val

145	Pro	Leu	Pro	Gly	Arg	150	Asp	Leu	Leu	Leu	Val	155	Ala	Arg	Gly	Leu	Ala	160	Gly
						165					170							175	
Lys	Leu	Ser	Ala	Gly	Val	Pro	Pro	Leu	Ala	Ile	Arg	Gln	Leu	Ala	Gly				
						180												190	
Ser	Gln	Pro	Phe	Ser	Ser	Glu	Gly	Leu	Gly	Arg	Leu	Val	Val	Gly	Asp				
						195												205	
Phe	Ser	Asp	Tyr	Asn	Asn	Ser	Tyr	Val	Gly	Ala	Phe	Ala	Asp	Ala	Arg				
						210												220	
Ser	Ala	Tyr	Phe	Val	Phe	Arg	Arg	Arg	Gly	Ala	Arg	Ala	Gln	Ala	Glu				
						225												235	
Tyr	Arg	Ser	Tyr	Val	Ala	Arg	Val	Cys	Leu	Gly	Asp	Thr	Asn	Leu	Tyr				
						245												255	
Ser	Tyr	Val	Glu	Val	Pro	Leu	Ala	Cys	Gln	Gly	Gln	Gly	Leu	Ile	Gln				
						260												270	
Ala	Ala	Phe	Leu	Ala	Pro	Gly	Thr	Leu	Leu	Gly	Val	Phe	Ala	Ala	Gly				
						275												285	
Pro	Arg	Gly	Thr	Gln	Ala	Ala	Leu	Cys	Ala	Phe	Pro	Met	Val	Glu	Leu				
						290												300	
Gly	Ala	Ser	Met	Glu	Gln	Ala	Arg	Arg	Leu	Cys	Tyr	Thr	Ala	Gly	Gly				
						310												320	
Arg	Gly	Pro	Ser	Gly	Ala	Glu	Glu	Ala	Thr	Val	Glu	Tyr	Gly	Val	Thr				
						325												335	
Ser	Arg	Cys	Val	Thr	Leu	Pro	Leu	Asp	Ser	Pro	Glu	Ser	Tyr	Pro	Cys				
						340												350	
Gly	Asp	Glu	His	Thr	Pro	Ser	Pro	Ile	Ala	Gly	Arg	Gln	Pro	Leu	Glu				
						355												365	
Val	Gln	Pro	Leu	Leu	Lys	Leu	Gly	Gln	Pro	Val	Ser	Ala	Val	Ala	Ala				
						370												380	
Leu	Gln	Ala	Asp	Gly	His	Met	Ile	Ala	Phe	Leu	Gly	Asp	Thr	Gln	Gly				
						385												400	
Gln	Leu	Tyr	Lys	Val	Phe	Leu	His	Gly	Ser	Gln	Gly	Gln	Val	Tyr	His				
						405												415	
Ser	Gln	Gln	Val	Gly	Pro	Pro	Gly	Ser	Ala	Ile	Ser	Pro	Asp	Leu	Leu				
						420												430	
Leu	Asp	Ser	Ser	Gly	Ser	His	Leu	Tyr	Val	Leu	Thr	Ala	His	Gln	Val				
						435												445	
Asp	Arg	Ile	Pro	Val	Ala	Ala	Cys	Pro	Gln	Phe	Pro	Asp	Cys	Ala	Ser				
						450												460	
Cys	Leu	Gln	Ala	Gln	Asp	Pro	Leu	Cys	Gly	Trp	Cys	Val	Leu	Gln	Gly				
						465												475	
Arg	Cys	Thr	Arg	Lys	Gly	Gln	Cys	Gly	Arg	Ala	Gly	Gln	Leu	Asn	Gln				
						485												490	
Trp	Leu	Trp	Ser	Tyr	Glu	Glu	Asp	Ser	His	Cys	Leu	His	Ile	Gln	Ser				
						500												510	
Leu	Leu	Pro	Gly	His	His	Pro	Arg	Gln	Glu	Gln	Gly	Gln	Val	Thr	Leu				
						515												525	
Ser	Val	Pro	Arg	Leu	Pro	Ile	Leu	Asp	Ala	Asp	Glu	Tyr	Phe	His	Cys				
						530												540	
Ala	Phe	Gly	Asp	Tyr	Asp	Ser	Leu	Ala	His	Val	Glu	Gly	Pro	His	Val				
						545												555	
Ala	Cys	Val	Thr	Pro	Pro	Gln	Asp	Gln	Val	Leu	Asn	Pro	Pro	Gly					
						555												570	
Thr	Asp	His	Val	Thr	Val	Pro	Leu	Ala	Leu	Met	Phe	Glu	Asp	Val	Thr				
						580												585	
																		590	

Val Ala Ala Thr Asn Phe Ser Phe Tyr Asp Cys Ser Ala Val Gln Ala
 595
 Leu Glu Ala Ala Pro Val Leu Pro Gln Gly Leu Pro Ala Ser Phe
 610
 His Cys Trp Leu Glu Leu Pro Gly Glu Leu Arg Gly Leu Pro Ala Thr
 625
 Leu Glu Glu Thr Ala Gly Asp Ser Gly Leu Ile His Cys Gln Ala His
 645
 Gln Arg Glu Leu Pro Val Pro Ile Tyr Val Thr Gln Gly Glu Ala Gln
 660
 Arg Leu Asp Asn Thr His Ala Leu Tyr Gly Glu Pro Glu Gly Ser Gln
 675
 Ala Gly Gly Ala Gly Trp Val Ala Asp Arg Arg Ser Ala His Cys
 690
 Leu Thr Leu Pro Ser Asp Pro Val Arg Leu Arg His Gly Pro Pro Gly
 705
 Leu Gln Pro Leu Pro Ser Gly Gln Gln Glu Pro Gly Leu Pro Val Thr
 725
 Ser Pro Ala Pro Gly Pro Gln Thr Pro Ala Ala Arg Pro Gly Trp Ala
 740
 Gly Trp Leu Ala Gly His Pro Ala Leu Gln Ser Gly Ala Trp Val Arg
 755
 Gly Thr Pro Ser Ala Ile Ile Cys Leu Leu Gln Val Glu Pro Leu Thr
 770
 Gly Pro Pro Glu Gly Gly Leu Ala Leu Thr Ile Leu Gly Ser Asn Leu
 785
 Gly Arg Ala Phe Ala Asp Val Gln Tyr Ala Asp Pro Val Leu Leu Ser
 805
 Leu Ser Pro Arg Trp Gly Pro Gln Ala Gly Gly Thr Gln Leu Thr Ile
 820
 Arg Gly Gln His Leu Gln Thr Gly Gly Asn Thr Ser Ala Phe Val Gly
 835
 Gly Gln Pro Cys Pro Met Gly Gly Arg Leu Ile Arg Val Arg Gly Thr
 850
 Gly Leu Asp Val Val Gln Arg Pro Leu Leu Ser Val Trp Leu Glu Ala
 865
 Asp Ala Glu Val Gln Ala Ser Arg Ala Gln Pro Gln Asp Pro Gln Pro
 885
 Arg Arg Ser Cys Gly Ala Pro Ala Ala Asp Pro Gln Ala Cys Ile Gln
 900
 Leu Gly Gly Gly Leu Leu Gln Arg Thr Ala Glu Pro Ser Ser Leu His
 915
 Leu Trp Ser Ala Leu Asn Ala Pro Gln Cys Ser Thr Val Cys Ser Val
 930
 Asn Ser Ser Ser Leu Leu Leu Cys Arg Ser Pro Ala Val Pro Asp Arg
 945
 Ala His Pro Gln Arg Val Phe Phe Thr Leu Asp Asn Val Gln Val Asp
 965
 Phe Ala Ser Ala Ser Gly Gly Gln Gly Phe Leu Tyr Gln Pro Asn Pro
 980
 Arg Leu Ala Pro Leu Ser Arg Glu Gly Pro Ala Arg Pro Tyr Arg Leu
 995
 Lys Pro Gly His Val Leu Asp Val Glu Gly Glu Gly Leu Asn Leu Gly
 1010
 Ile Ser Lys Glu Glu Val Arg Val His Ile Gly Arg Gly Glu Cys Leu
 1015

1025		1030		1035		1040
Val Lys Thr Leu	Thr Arg Thr His	Leu Tyr Cys Glu	Pro Pro Ala His			
	1045	1050	1055			
Ala Pro Gln Pro	Ala Asn Gly Ser Gly	Leu Pro Gln Phe	Val Val Gln			
	1060	1065	1070			
Met Gly Asn Val	Gln Leu Ala Leu	Gly Pro Val Gln	Tyr Glu Ala Glu			
	1075	1080	1085			
Pro Pro Leu Ser	Ala Phe Pro Val	Glu Ala Gln Ala	Gly Val Gly Met			
	1090	1095	1100			
Gly Ala Ala Val	Leu Ile Ala Ala	Val Leu Leu Thr	Leu Met Tyr			
	1105	1110	1115			
Arg His Lys Ser	Lys Gln Ala Leu	Arg Asp Tyr Gln	Lys Val Leu Val			
	1125	1130	1135			
Gln Leu Glu Ser	Leu Glu Thr Gly	Val Gly Asp Gln	Cys Arg Lys Glu			
	1140	1145	1150			
Phe Thr Asp Leu	Met Thr Glu Met	Thr Asp Leu Ser	Ser Asp Leu Glu			
	1155	1160	1165			
Gly Ser Gly Ile	Pro Phe Leu Asp	Tyr Arg Thr Tyr	Ala Glu Arg Ala			
	1170	1175	1180			
Phe Phe Pro Gly	His Gly Gly Cys	Pro Leu Gln Pro	Lys Pro Glu Gly			
	1185	1190	1195			
Pro Gly Glu Asp	Gly His Cys Ala	Thr Val Arg Gln	Gly Leu Thr Gln			
	1205	1210	1215			
Leu Ser Asn Leu	Leu Asn Ser Lys	Leu Phe Leu Thr	Val Arg Ala			
	1220	1225	1230			
Val Trp Arg Glu	Cys Pro Val Gly	Lys Glu Val Gly	Leu Asn Tyr			
	1235	1240	1245			
Trp Pro Glu Thr	Lys Val Gly Glu	Glu Thr Glu Thr	Met Val Glu Lys			
	1250	1255	1260			
Leu Leu Thr Asn	Trp Leu Ser Ile	Cys Leu Tyr Ala	Phe Leu Arg Glu			
	1265	1270	1275			
Val Ala Gly Glu	Pro Leu Tyr Met	Leu Phe Arg Ala	Ile Gln Tyr Gln			
	1285	1290	1295			
Val Asp Lys Gly	Pro Val Asp Ala	Val Thr Gly Lys	Ala Lys Arg Thr			
	1300	1305	1310			
Leu Asn Asp Ser	Arg Leu Leu Arg	Glu Asp Val Glu	Phe Gln Pro Leu			
	1315	1320	1325			
Thr Leu Met Val	Leu Val Gly Pro	Gly Ala Gly Gly	Ala Ala Gly Ser			
	1330	1335	1340			
Ser Glu Met Gln	Arg Val Pro Ala	Arg Val Leu Asp	Thr Asp Thr Ile			
	1345	1350	1355			
Thr Gln Val Lys	Glu Lys Val Leu	Asp Gln Val Tyr	Lys Gly Thr Pro			
	1365	1370	1375			
Phe Ser Gln Arg	Pro Ser Val His	Ala Leu Asp Leu	Gly Glu Arg Ala			
	1380	1385	1390			
Ser Pro Ala His	Pro Pro Gln Gly	Pro Phe Pro Thr	Pro Pro Ala Pro			
	1395	1400	1405			
Gly Ala Pro Gln	Leu Cys Leu Thr	Met Asn Ile Pro	Thr Leu Glu Asp			
	1410	1415	1420			
Gly Glu Glu Gly	Gly Val Cys Leu	Trp His Leu Val	Lys Ala Thr Glu			
	1425	1430	1435			
Glu Pro Glu Gly	Ala Lys Val Arg	Cys Ser Ser Leu	Arg Glu Arg Glu			
	1445	1450	1455			
Pro Ala Arg Ala	Lys Ala Ile Pro	Glu Ile Tyr Leu	Thr Arg Leu Leu			
	1460	1465	1470			

Ser Met Lys Val Gly Ala Ala Trp Val Ala Gly Pro Glu Arg Arg Leu
 1475 1480 1485
 Ser Gln Gly Pro Arg Pro Ser Gln Gly Val Gly Gly Ala Gly Ala Ala
 1490 1495 1500
 Ser Ala Val Asp Gly Pro His Thr Leu Pro Ser Thr Gln Pro Leu Ser
 1505 1510 1515 1520
 Pro Ala Ser Gln Gly Thr Leu Gln Lys Phe Val Asp Asp Thr Phe Gln
 1525 1530 1535
 Ala Ile Leu Ser Val Asn Arg Pro Ile Pro Ile Ala Val Lys Tyr Leu
 1540 1545 1550
 Phe Asp Leu Leu Asp Glu Leu Ala Glu Lys His Gly Ile Glu Asp Pro
 1555 1560 1565
 Gly Thr Leu His Ile Trp Lys Thr Asn Ser Leu Leu Arg Phe Trp
 1570 1575 1580
 Val Asn Ala Leu Lys Asn Pro Gln Leu Ile Phe Asp Val Arg Val Ser
 1585 1590 1595 1600
 Asp Asn Val Asp Ala Ile Leu Ala Val Ile Ala Gln Thr Phe Ile Asp
 1605 1610 1615
 Ser Cys Thr Thr Ser Glu His Lys Val Gly Arg Val Arg Ala Val Pro
 1620 1625 1630
 Ala Ala Ala Ala Gly Arg Gly Leu Arg Arg Lys Gly Leu Trp Gly Lys
 1635 1640 1645
 Pro Arg Gly Leu Cys Thr Glu Leu Trp Val Gly Ser Gly Ser Ile Met
 1650 1655 1660
 Gly Ala Pro Ser Pro Pro Ser Ser Cys Leu Ala Pro Pro Leu Pro Pro
 1665 1670 1675 1680
 Glu Gln Asp Ser Pro Val Asn Lys Leu Leu Tyr Ala Arg Glu Ile Pro
 1685 1690 1695
 Arg Tyr Lys Gln Met Val Glu Arg Tyr Tyr Ala Asp Ile Arg Gln Ser
 1700 1705 1710
 Ser Pro Ala Ser Tyr Gln Glu Met Asn Ser Ala Leu Ala Glu Leu Ser
 1715 1720 1725
 Gly Asn Tyr Thr Ser Ala Pro His Cys Leu Glu Ala Leu Gln Glu Leu
 1730 1735 1740
 Tyr Asn His Ile His Arg Tyr Tyr Asp Gln Ile Ser Ala Leu Glu
 1745 1750 1755 1760
 Glu Asp Pro Val Gly Gln Lys Leu Gln Leu Ala Cys Arg Leu Gln Gln
 1765 1770 1775
 Val Ala Ala Leu Val Glu Asn Lys Val Thr Asp Leu
 1780 1785

<210> 5

<211> 5892

<212> DNA

<213> HOMO SAPIEN

<400> 5

gcgcacgccc	ggatggctct	tcgcgcgcgcg	ggcggcgccac	cccttagcgg	cccggccgcc	60
gcgcacgccc	cccgcgcgtt	ccagacgcgc	ccgcgggtgcc	cggtgccgct	gctgttgctg	120
ctgtctcctgg	ggcggcgctg	ggcggcgccc	ctggagatcc	agcgtcggtt	cccctcgccc	180
acgcccaccca	acaaacttcgc	ctggagcggc	cgcgccggga	ccgtgtacct	ggcggccgtc	240
aaccgcctct	atcagctgtc	gggcgccaac	ctgagcttgg	agccggaggc	ggcctgtggc	300
ccggtgcccc	acagcccgcct	gtgtcaagct	ccgcagctgc	cgcaggcctc	gtgcgagcac	360
ccgcggcgcc	tcaaggacaa	ctacaacaag	atcctgcagc	tggaccccgc	ccaggccctg	420
gtagctcgtg	gcgggtccat	ctaccagggc	ttctgccagc	tgccgcccgc	gggtaacatc	480

tcggcgcgtg	cogtgcgctt	cccgcccgcc	gcgcgcgcgc	ccgagccgct	cacggtgttc	540
cccgcatgct	tgaagctggc	ggccaaccac	ccgaacgcgt	ccaccgtggg	gctagtctgt	600
ctcccccgcg	ggggcgccgg	gggcagccgc	ctgctcgtgg	ggcccacgta	caccggttac	660
ggcagctctg	tcttcccgcg	caaccgcagc	ctggaggacc	accgcttcga	gaacacgccc	720
gagatcgcca	tccgtccctc	ggacacgcgc	ggcgacctgt	ccaagctctt	caccctcgac	780
ctcaaccctc	ccgacgacaa	catcctcaag	atcaagcaga	ggccacacag	gcagcaaaag	840
ctcgctctcg	tgagcgccct	ctcgcaaccg	tccgaccgcg	cgccgggtgc	acagtctcac	900
gcgtacacgc	cgctcaaacg	cgaggcgccg	gcggggcgca	ggcgcggagc		960
ctcttcggcg	gcattctgct	gccccaacgg	ccggcgccgg	acgccaagaa	gctcacccag	1020
tctacacttc	agtctggcct	tcagtgccgc	ggcgccgcgc	ggcggcgcca	ctctctacag	1080
cgctcgttgt	cggtcttccc	agccccggag	cggtctcttg	ctgtcttcga	gcggccccag	1140
gggttccccg	ggcccgccgc	tgctccggcg	gcactctgcy	cttctcgctt	cgccgcagtg	1200
cgagcccgca	ctcgagctgc	gcgcaccgcc	tgcttcgtgg	aacggcgccc	cgacgtgggtg	1260
gcggtgtctg	acagcgttgt	cgaggcgacg	ggacccgctc	gcgagcgcaa	gctcaaacat	1320
cagctccccc	cagagcagct	ggactgttga	gctgctcacc	tcagcacccc	gctgtccate	1380
ctcgaccctc	tgaaggccac	cccgctgttc	cgcgccccgc	gcctcacctc	cggtggcgctg	1440
gccagcgtca	acaactacac	agcgctcttc	ctgggacagg	tcaacgggag	gcttctcaag	1500
atcaaccctc	acagagacat	gcaggtgggt	agcagggcgg	tggtgactgt	ggcctatggg	1560
gagccccgtg	accatgtcat	gcagtttgac	ccagcagact	ccggttaact	ttacctgatg	1620
acgtcccaac	agatggccag	ggtagagtcg	gcgcctcgca	acgtgctgca	cactctgtgg	1680
gtctcgctgg	gtgcggcgga	cgctactctg	ggctggtgtg	ccctggagac	gcggtgcacc	1740
ttagcagcag	actgcaccaa	ttccagccag	cagcatttct	ggaccagctc	cagcgagggc	1800
cccagccgct	gtctctccat	gaccgtctct	cttcccgaga	tcgatgtgcy	ccaggagatc	1860
ccaggactga	tccttcagat	ctcgggcagc	ctgccacgcc	tcagtgactgc	ggagatggcc	1920
tgtagactat	gggaacaact	ccgcactgtg	gctcgggtcc	caggccctgc	ctttgtgtcc	1980
cagatttgcct	actgcaacct	ctgcggagg	gaccagtttc	gcctcttccc	ccccaaaccag	2040
gaccacgtga	ctgttgagat	gtctgtgagg	gtcaatgggc	ggaacatcgt	caaggccaat	2100
ttaccatctc	acgactgcag	ccgcactgca	caagtgtacc	cccacacagc	ctgtaccagc	2160
tgctctgcgg	cacagtggcc	ctgtttcttg	tgacgcccag	agcactctct	tgtttccaac	2220
cagttctcct	gcgagccctc	accaaaaccc	acgagccctc	aggactgccc	ccggaccctg	2280
ctctcacccc	tggcacccgt	gcttaccggt	ggctcccaga	acatctctgt	gctcttgccc	2340
aacactgcct	ttttccaggg	tgacgcccgt	gagtgtagtt	ttgggctgga	ggagatcttc	2400
gaggctgtgt	gggtgaatga	gtctgttgta	cgctgtgacc	aggtgggtgt	gcacacgacc	2460
cggaaagacc	aggtgttccc	gctcagccct	caactaaagg	ggcggccagc	ccgatctctg	2520
gacagccctg	agcccatgac	agtcatgggtc	tataactgtg	ccatgggcag	ccccgactgt	2580
ttccactgct	tgggcccgga	agacctgggt	caactgtgcy	tggtgagtag	tggtctgccg	2640
ctcggggggc	ctctgcagcc	catggctggc	acctgccccg	ccccgagat	cggcgcatgt	2700
gagccccctg	gtggcccggt	ggacggtggg	acctgcctga	ccatccgagg	aagggaacctg	2760
ggccggcgcc	tcagtgactgt	ggcccacggc	gtgtggattg	gtggtgtggc	ctgtgagcca	2820
ctccttgaca	gatacacggt	gtcgaggagg	atcgtgtgtg	tcacagggcg	agccccagga	2880
ccctctctag	gtgtggtgac	cgtaaacgcc	tctaaggagg	gcaagtcccc	ggaccgcttc	2940
tctcactgct	tgccctcggt	ccactccctg	gagcctaaca	tgggccccaa	ggccgggggc	3000
accaggatca	ccctccatgg	gaatgacctc	catgtagget	ccgagctcca	ggctctggtg	3060
aacgacacag	acccttgcaac	ggagctgatg	cgcacagata	ccagcatcgt	gctgaacctg	3120
cccgaggggc	ccctgcggcg	tcgggtgctt	gtgtgtgtgc	gctcagagcg	tcggggctgc	3180
gtcaccggca	accctcacctt	ctggtacatg	cagaaccggg	tcactcaggc	catcagctcc	3240
cgcccgagcc	ctgtcagtag	cggcaggacc	atccagatgg	ctggtgagcg	tttccacatg	3300
gtcgagaatg	tgtccatggc	gctccacacc	atctggccggg	agccccagtg	cctcagaggtt	3360
ctcactccca	ccctcatcac	ctgcccgttc	ccggggggcc	tgaagcaacc	atcagcgcca	3420
gtggactctc	tctcaaatgg	gcgggctcac	gcgagcaggg	tggtgtgggc	tgaaggagcta	3480
ctgtcccccg	aggagggcaca	gcggggcagc	aggttccgct	tggaactacct	cccccaacca	3540
cagttctctca	ggcccaagag	ggagaagtag	atcaagcacc	accccgggga	gcctctcacc	3600
ctcgttatcc	acaaggagca	ggacagcctg	gggctccaga	gtccagagta	cggggtcaag	3660
ataggccaag	taagctcgga	catccagatt	gtctctgaca	gatacatcca	ctctctgggtc	3720
aacgagctcc	tgggcggcgc	cgttggggcag	ctgcccataca	caatccaggt	agggaaacttc	3780

aaccagacca	tcgccacact	gcagctgggg	ggcagcgaga	cggccatcat	cgtgtccatc	3840
gtcatctgca	gcgtcctgct	gctgctctcc	gtggtggccc	tgttcgtctt	ctgtaccaag	3900
agccgcacgt	ctgagcgtta	ctggcgagaag	acgctgctgc	agatggagga	gatggaaatc	3960
cagatccgag	aggaataatccg	caaaggcttc	gctgagctgc	agacagacat	gacagatctg	4020
accaaggcag	tgaaccgcag	ccaggcgatc	cccttctctg	agtataagca	cttcgtgacc	4080
cgcaactttt	tcaccaagtg	ttctctccctt	tatgaagagc	gttactgtgt	gccctcccag	4140
accctcaact	cccaggggcag	ctcccaggga	caggaaaacc	acccactgct	gggagagtgg	4200
aagatctcct	agagctgcgc	gcccacaact	gaagaggga	ttagcgtgtt	ctcctcaacta	4260
ctcaacaaca	agcacttctt	catcgtcttt	gtccacgcgc	tggagcagca	gaaggacttt	4320
gcggtgcgcg	acaggttgacg	cttggcctcg	ctgctgacca	tcgcgctgca	cggaagctgt	4380
gagtgactaca	ccagcatcat	gaaggagctg	ctggtggacc	tcattgacgc	ctcggccgcc	4440
aagaacccca	agctcatgct	cgggcgaca	gagtcgtggt	tggagaagat	gctcaccaac	4500
tggatgtcca	tcctcatgta	cagctgtctg	cgggagacgg	tgggggagct	attcttctctg	4560
ctgctgtgtg	ccatcaagca	gcaaatcaac	aaagggtcca	tcgacgccat	cacaggcaag	4620
gcccgttaca	cactcaatga	ggagtggctg	ctcggggaga	acatcgagcg	caagccccgg	4680
aaactgaacg	tgtcttctca	gggctgtggc	atggactcgc	tgaagctgcy	ggccatggag	4740
accgacacgc	tgacacaggt	caagggaag	alcctggagg	ccctctgcaa	gaatgtgcc	4800
tactccagct	ggccgcgtgc	agaggacgtc	gaccttgagt	ggttcgccc	cagcacacag	4860
agctacatcc	ttcgggaact	ggacgacacc	tcagtggagg	aagacggccc	caagaagctt	4920
aaacagctgg	ccattataca	gatccctgaa	ggctgcctcc	tggcatgag	tctcatagac	4980
aagaaggaca	acacactggg	cgagtgaaa	gacttggaca	cagagaagt	ttctcattg	5040
gtgctgccta	cgacagagct	ggcgagccc	aagaagcttc	accggagag	catcgacaag	5100
aaagtgctcc	cggaaatcta	ctgacccgc	ctgctctcca	caaaggccac	gttcagaaag	5160
ttcttgatg	acctgttcaa	ggccattctg	agtatccgtg	aagacaagcc	cccactggct	5220
gtcaagtact	ttttcgactt	ctctgaggag	cagcgtgaga	agaagggaat	ctccgacccc	5280
gacacacctc	acatctggaa	gaccaacagc	cttcctctcc	ggttctgggt	gaacattctg	5340
aagaacccccc	agtttgtctt	tgacatcgac	aagaacagacc	acatcgagc	ctgcctttca	5400
gtcatcgcgc	agggcttcat	cgacgcctgc	tccatctctg	acctgcagct	gggcaaggat	5460
tgcacaacca	acaagctctt	ctacgcacaag	gagattctctg	agtaaccggaa	gatcgtgacg	5520
cgctactaca	acgagatcca	ggacatgacg	ccgctcagcg	agcaagagat	gaatgcccat	5580
ctggcccgagg	agtcagaggaa	ataccagaat	gagttcaaca	ccaatgtggc	catggcagag	5640
attttttaggt	cgcccaagag	gtatcggccg	cagatcatgg	ccgcgttgga	ggccaacccc	5700
acggcccgga	ggacacaact	gcagcacaag	tttgagcagg	tggttggctt	gatggaggac	5760
aaactctacg	agtgctaacg	tgaggctctga	gacacatgga	gagttgtcta	ggctgctgct	5820
gggagaatg	gacgccact	gggcctcaac	ttgatcttct	accccggtcc	tgtgactcag	5880
actgggaat	ac					5920

<210> 6
 <211> 1925
 <212> PRT
 <213> HOMO SAPIEN

<400> 6
 Met Ala Leu Arg Ala Ala Gly Gly Ala Pro Phe Ser Gly Pro Ala Ala
 1 5 10 15
 Ala Ala Ser Pro Pro Phe Gln Thr Pro Pro Arg Cys Pro Val Pro
 20 25 30
 Leu Leu Leu Leu Leu Leu Gly Ala Ala Arg Ala Gly Ala Leu Glu
 35 40 45
 Ile Gln Arg Arg Phe Pro Ser Pro Thr Pro Thr Asn Asn Phe Ala Leu
 50 55 60
 Asp Gly Ala Ala Gly Thr Val Tyr Leu Ala Ala Val Asn Arg Leu Tyr
 65 70 75 80

Gln Leu Ser Gly Ala Asn Leu Ser Leu Glu Ala Glu Ala Ala Val Gly
 85 90 95
 Pro Val Pro Asp Ser Pro Leu Cys His Ala Pro Gln Leu Pro Gln Ala
 100 105
 Ser Cys Glu His Pro Arg Arg Leu Thr Asp Asn Tyr Asn Lys Ile Leu
 115 120 125
 Gln Leu Asp Pro Gly Gln Gly Leu Val Val Val Cys Gly Ser Ile Tyr
 130 135 140
 Gln Gly Phe Cys Gln Leu Arg Arg Arg Gly Asn Ile Ser Ala Val Ala
 145 150 155
 Val Arg Phe Pro Pro Ala Ala Pro Pro Ala Glu Pro Val Thr Val Phe
 165 170 175
 Pro Ser Met Leu Asn Val Ala Ala Asn His Pro Asn Ala Ser Thr Val
 180 185 190
 Gly Leu Val Leu Pro Pro Ala Ala Gly Ala Gly Gly Ser Arg Leu Leu
 195 200 205
 Val Gly Ala Thr Tyr Thr Gly Tyr Gly Ser Ser Phe Phe Pro Arg Asn
 210 215 220
 Arg Ser Leu Glu Asp His Arg Phe Glu Asn Thr Pro Glu Ile Ala Ile
 225 230 235 240
 Arg Ser Leu Asp Thr Arg Gly Asp Leu Ala Lys Leu Phe Thr Phe Asp
 245 250 255
 Leu Asn Pro Ser Asp Asp Asn Ile Leu Lys Ile Lys Gln Gly Ala Lys
 260 265 270
 Glu Gln His Lys Leu Gly Phe Val Ser Ala Phe Leu His Pro Ser Asp
 275 280 285
 Pro Pro Pro Gly Ala Gln Ser Tyr Ala Tyr Leu Ala Leu Asn Ser Glu
 290 295 300
 Ala Arg Ala Gly Asp Lys Glu Ser Gln Ala Arg Ser Leu Leu Ala Arg
 305 310 315 320
 Ile Cys Leu Pro His Gly Ala Gly Gly Asp Ala Lys Lys Leu Thr Glu
 325 330 335
 Ser Tyr Ile Gln Leu Gly Leu Gln Cys Ala Gly Gly Ala Gly Arg Gly
 340 345 350
 Asp Leu Tyr Ser Arg Leu Val Ser Val Phe Pro Ala Arg Glu Arg Leu
 355 360 365
 Phe Ala Val Phe Glu Arg Pro Gln Gly Ser Pro Ala Ala Arg Ala Ala
 370 375 380
 Pro Ala Ala Leu Cys Ala Phe Arg Phe Ala Asp Val Arg Ala Ala Ile
 385 390 395 400
 Arg Ala Ala Arg Thr Ala Cys Phe Val Glu Pro Ala Pro Asp Val Val
 405 410 415
 Ala Val Leu Asp Ser Val Val Gln Gly Thr Gly Pro Ala Cys Glu Arg
 420 425 430
 Lys Leu Asn Ile Gln Leu Gln Pro Glu Gln Leu Asp Cys Gly Ala Ala
 435 440 445
 His Leu Gln His Pro Leu Ser Ile Leu Gln Pro Leu Lys Ala Thr Pro
 450 455 460
 Val Phe Arg Ala Pro Gly Leu Thr Ser Val Ala Val Ala Ser Val Asn
 465 470 475 480
 Asn Tyr Thr Ala Val Phe Leu Gly Thr Val Asn Gly Arg Leu Leu Lys
 485 490 495
 Ile Asn Leu Asn Glu Ser Met Gln Val Val Ser Arg Arg Val Val Thr
 500 505 510
 Val Ala Tyr Gly Glu Pro Val His His Val Met Gln Phe Asp Pro Ala

Ays	530	515				520				525						
		Ser	Gly	Tyr	Leu	Tyr	Leu	Met	Thr	Ser	His	Gln	Met	Ala	Arg	Val
Lys	Val	Ala	Ala	Cys	Asn	Val	His	Ser	Thr	Cys	555	Gly	Asp	Cys	Val	Gly
545	Ala	Ala	Asp	Ala	Tyr	Cys	Gly	Trp	Cys	Ala	Leu	Glu	Thr	Arg	Cys	Thr
Leu	Gln	Gln	Asp	Cys	Thr	Asn	Ser	Ser	Gln	Gln	His	Phe	Trp	Thr	Ser	Thr
Ala	Ser	Glu	Gly	Pro	Ser	Arg	Cys	Pro	Ala	Met	Thr	Val	Leu	Pro	Ser	Thr
Glu	Ile	Asp	Val	Arg	Gln	Glu	Tyr	Pro	Gly	Met	Ile	Leu	Gln	Ile	Ser	Thr
Gly	Ser	Leu	Pro	Ser	Leu	Ser	Gly	Met	Glu	Met	Ala	Cys	Asp	Tyr	Gly	Thr
625	Asn	Asn	Ile	Arg	Thr	Val	Ala	Arg	Val	Pro	Gly	Pro	Ala	Phe	Gly	His
Asn	Ile	Ala	Tyr	Cys	Asn	Leu	Leu	Pro	Arg	Asp	Gln	Phe	Pro	Pro	Phe	Thr
Pro	Pro	Asn	Gln	Asp	His	Val	Thr	Thr	Val	Glu	Met	Ser	Val	Arg	Val	Asn
Gly	Arg	Asn	Ile	Val	Lys	Ala	Asn	Phe	Thr	Ile	Tyr	Asp	Cys	Ser	Arg	Thr
Thr	Ala	Gln	Val	Tyr	Pro	His	Thr	Ala	Cys	Thr	Ser	Cys	Leu	Ser	Ala	Thr
Gln	Trp	Pro	Cys	Phe	Trp	Cys	Ser	Gln	Gln	His	Ser	Cys	Val	Ser	Asn	Thr
Gln	Ser	Arg	Cys	Glu	Ala	Ser	Pro	Asn	Pro	Thr	Ser	Pro	Gln	Asp	Cys	Thr
Pro	Arg	Thr	Leu	Leu	Ser	Pro	Leu	Ala	Pro	Val	Pro	Thr	Gly	Gly	Ser	Thr
Gln	Asn	Ile	Leu	Val	Pro	Leu	Ala	Asn	Thr	Ala	Phe	Phe	Gln	Gly	Ala	Thr
Ala	Leu	Glu	Cys	Ser	Phe	Gly	Leu	Glu	Glu	Ile	Phe	Glu	Ala	Val	Trp	Thr
785	Asn	Glu	Ser	Val	Val	Arg	Cys	Asp	Gln	Val	Val	Leu	His	Thr	Thr	Thr
Arg	Lys	Ser	Gln	Val	Phe	Pro	Leu	Ser	Leu	Gln	Leu	Lys	Gly	Arg	Pro	Thr
Ala	Arg	Phe	Leu	Asp	Ser	Pro	Glu	Pro	Met	Thr	Val	Met	Val	Tyr	Asn	Thr
Cys	Ala	Met	Gly	Ser	Pro	Asp	Cys	Ser	Gln	Cys	Leu	Gly	Arg	Glu	Asp	Thr
Leu	Gly	His	Leu	Cys	Val	Trp	Ser	Asp	Gly	Cys	Arg	Leu	Arg	Gly	Pro	Thr
865	Leu	Gln	Pro	Met	Ala	Gly	Thr	Cys	Pro	Ala	Pro	Glu	Ile	Arg	Ala	Ile
Glu	Pro	Leu	Ser	Gly	Pro	Leu	Asp	Gly	Gly	Thr	Leu	Leu	Thr	Ile	Arg	Thr
Gly	Arg	Asn	Leu	Gly	Arg	Arg	Leu	Ser	Asp	Val	Ala	His	Gly	Val	Trp	Thr
Ile	Gly	Gly	Val	Ala	Cys	Glu	Pro	Leu	Pro	Asp	Arg	Tyr	Thr	Val	Ser	Thr
Glu	Glu	Ile	Val	Cys	Val	Thr	Gly	Pro	Ala	Pro	Gly	Pro	Leu	Ser	Gly	Thr
945					950					955						

Val Val Thr Val Asn Ala Ser Lys Glu Gly Lys Ser Arg Asp Arg Phe
 965 970 975
 Ser Tyr Val Leu Pro Leu Val His Ser Leu Glu Pro Thr Met Gly Pro
 980 985 990
 Lys Ala Gly Gly Thr Arg Ile Thr Ile His Gly Asn Asp Leu His Val
 995 1000 1005
 Gly Ser Glu Leu Gln Val Leu Val Asn Asp Thr Asp Pro Cys Thr Glu
 1010 1015 1020
 Leu Met Arg Thr Asp Thr Ser Ile Ala Cys Thr Met Pro Glu Gly Ala
 1025 1030 1035 1040
 Leu Pro Ala Pro Val Pro Val Cys Val Arg Phe Glu Arg Arg Gly Cys
 1045 1050 1055
 Val His Gly Asn Leu Thr Phe Trp Tyr Met Gln Asn Pro Val Ile Thr
 1060 1065 1070
 Ala Ile Ser Pro Arg Arg Ser Pro Val Ser Gly Gly Arg Thr Ile Thr
 1075 1080 1085
 Val Ala Gly Glu Arg Phe His Met Val Gln Asn Val Ser Met Ala Val
 1090 1095 1100
 His His Ile Gly Arg Glu Pro Thr Leu Cys Lys Val Leu Asn Ser Thr
 1105 1110 1115 1120
 Leu Ile Thr Cys Pro Ser Pro Gly Ala Leu Ser Asn Ala Ser Ala Pro
 1125 1130 1135
 Val Asp Phe Phe Ile Asn Gly Arg Ala Tyr Ala Asp Glu Val Ala Val
 1140 1145 1150
 Ala Glu Glu Leu Leu Asp Pro Glu Glu Ala Gln Arg Gly Ser Arg Phe
 1155 1160 1165
 Arg Leu Asp Tyr Leu Pro Asn Pro Gln Phe Ser Thr Ala Lys Arg Glu
 1170 1175 1180
 Lys Trp Ile Lys His His Pro Gly Glu Pro Leu Thr Leu Val Ile His
 1185 1190 1195 1200
 Lys Glu Gln Asp Ser Leu Gly Leu Gln Ser His Glu Tyr Arg Val Lys
 1205 1210 1215
 Ile Gly Gln Val Ser Cys Asp Ile Gln Ile Val Ser Asp Arg Ile Ile
 1220 1225 1230
 His Cys Ser Val Asn Glu Ser Leu Gly Ala Ala Val Gly Gln Leu Pro
 1235 1240 1245
 Ile Thr Ile Gln Val Gly Asn Phe Asn Gln Thr Ile Ala Thr Leu Gln
 1250 1255 1260
 Leu Gly Gly Ser Glu Thr Ala Ile Ile Val Ser Ile Val Ile Cys Ser
 1265 1270 1275 1280
 Val Leu Leu Leu Leu Ser Val Val Ala Leu Phe Val Phe Cys Thr Lys
 1285 1290 1295
 Ser Arg Arg Ala Glu Arg Tyr Trp Gln Lys Thr Leu Leu Gln Met Glu
 1300 1305 1310
 Glu Met Glu Ser Gln Ile Arg Glu Glu Ile Arg Lys Gly Phe Ala Glu
 1315 1320 1325
 Leu Gln Thr Asp Met Thr Asp Leu Thr Lys Glu Leu Asn Arg Ser Gln
 1330 1335 1340
 Gly Ile Pro Phe Leu Glu Tyr Lys His Phe Val Thr Arg Thr Phe Phe
 1345 1350 1355 1360
 Pro Lys Cys Ser Ser Leu Tyr Glu Glu Arg Tyr Val Leu Pro Ser Gln
 1365 1370 1375
 Thr Leu Asn Ser Gln Gly Ser Ser Gln Ala Gln Glu Thr His Pro Leu
 1380 1385 1390
 Leu Gly Glu Trp Lys Ile Pro Glu Ser Cys Arg Pro Asn Met Glu Glu

1395	1400	1405
Gly Ile Ser Val Phe Ser Ser Leu Leu Asn Asn Lys His Phe Leu Ile		
1410	1415	1420
Val Phe Val His Ala Leu Glu Gln Gln Lys Asp Phe Ala Val Arg Asp		
1425	1430	1435
Arg Cys Ser Leu Ala Ser Leu Leu Thr Ile Ala Leu His Gly Lys Leu		
1445	1450	1455
Glu Tyr Tyr Thr Ser Ile Met Lys Glu Leu Leu Val Asp Leu Ile Asp		
1460	1465	1470
Ala Ser Ala Ala Lys Asn Pro Lys Leu Met Leu Arg Arg Thr Glu Ser		
1475	1480	1485
Val Val Glu Lys Met Leu Thr Asn Trp Met Ser Ile Cys Met Tyr Ser		
1490	1495	1500
Cys Leu Arg Glu Thr Val Gly Glu Pro Phe Phe Leu Leu Leu Cys Ala		
1505	1510	1515
Ile Lys Gln Gln Ile Asn Lys Gly Ser Ile Asp Ala Ile Thr Gly Lys		
1525	1530	1535
Ala Arg Tyr Thr Leu Asn Glu Glu Trp Leu Leu Arg Glu Asn Ile Glu		
1540	1545	1550
Ala Lys Pro Arg Asn Leu Asn Val Ser Phe Gln Gly Cys Gly Met Asp		
1555	1560	1565
Ser Leu Ser Val Arg Ala Met Asp Thr Asp Thr Leu Thr Gln Val Lys		
1570	1575	1580
Glu Lys Ile Leu Glu Ala Phe Cys Lys Asn Val Pro Tyr Ser Gln Trp		
1585	1590	1595
Pro Arg Ala Glu Asp Val Asp Leu Glu Trp Phe Ala Ser Ser Thr Gln		
1605	1610	1615
Ser Tyr Ile Leu Arg Asp Leu Asp Asp Thr Ser Val Val Glu Asp Gly		
1620	1625	1630
Arg Lys Lys Leu Asn Thr Leu Ala His Tyr Lys Ile Pro Glu Gly Ala		
1635	1640	1645
Ser Leu Ala Met Ser Leu Ile Asp Lys Lys Asp Asn Thr Leu Gly Arg		
1650	1655	1660
Val Lys Asp Leu Asp Thr Glu Lys Tyr Phe His Leu Val Leu Pro Thr		
1665	1670	1675
Asp Glu Leu Ala Glu Pro Lys Lys Ser His Arg Gln Ser His Arg Lys		
1685	1690	1695
Lys Val Leu Pro Glu Ile Tyr Leu Thr Arg Leu Leu Ser Thr Lys Gly		
1700	1705	1710
Thr Leu Gln Lys Phe Leu Asp Asp Leu Phe Lys Ala Ile Leu Ser Ile		
1715	1720	1725
Arg Glu Asp Lys Pro Pro Leu Ala Val Lys Tyr Phe Phe Asp Phe Leu		
1730	1735	1740
Glu Glu Gln Ala Glu Lys Arg Gly Ile Ser Asp Pro Asp Thr Leu His		
1745	1750	1755
Ile Trp Lys Thr Asn Ser Leu Pro Leu Arg Phe Trp Val Asn Ile Leu		
1765	1770	1775
Lys Asn Pro Gln Phe Val Phe Asp Ile Asp Lys Thr Asp His Ile Asp		
1780	1785	1790
Ala Cys Leu Ser Val Ile Ala Gln Ala Phe Ile Asp Ala Cys Ser Ile		
1795	1800	1805
Ser Asp Leu Gln Leu Gly Lys Asp Ser Pro Thr Asn Lys Leu Leu Tyr		
1810	1815	1820
Ala Lys Glu Ile Pro Glu Tyr Arg Lys Ile Val Gln Arg Tyr Tyr Lys		
1825	1830	1835
		1840

Gln Ile Gln Asp Met Thr Pro Leu Ser Glu Gln Glu Met Asn Ala His
 1845 1850 1855
 Leu Ala Glu Glu Ser Arg Lys Tyr Gln Asn Glu Phe Asn Thr Asn Val
 1860 1865 1870
 Ala Met Ala Glu Ile Phe Arg Ser Pro Lys Arg Tyr Arg Pro Gln Ile
 1875 1880 1885
 Met Ala Ala Leu Glu Ala Asn Pro Thr Ala Arg Arg Thr Gln Leu Gln
 1890 1895 1900
 His Lys Phe Glu Gln Val Val Ala Leu Met Glu Asp Asn Ile Tyr Glu
 1905 1910 1915 1920
 Cys Tyr Ser Glu Ala
 1925

<210> 7
 <211> 601
 <212> DNA
 <213> HOMO SAPIEN

<400> 7
 caccagagtc cctgtggagt cctgtgggtca gtatcacagac tgcggcgagt gccttgggctc 60
 aggcgacccc cactgtggct ggtgtgtgct gcacaacact tgcacccgga aggcgagggtg 120
 tgagcggtcc aaggagcccc gcagggtttgc ctccggagatg aagcagtggtg tccgggtgac 180
 ggtccatccc aacaatatct ccgtctctca gtacaacgcg ctgctgggtcc tggagacgta 240
 caatgtcccc gagctgtcag ctggcgctcaa ctgcaccttt gaggacctgt cagagatgga 300
 tgggctggct gtgggcaatc agatccagtg ctactccctt gcagccaagg aggtgccccg 360
 gatcatcaca gagaatgggg accaccatgt cgtacagctt cagctcaaat caaaggagac 420
 cggcatgacc ttgccagca ccagctttgt ctctacaat tgcagcgtcc acaatctgtg 480
 cctgtctctgc gtggagagtc cataccgctg ccactgggtg aaataccgcg atgtctgcac 540
 ccatgacccc aagacctgct ccttccagga aggcgagtg aagctgccc aggtagggtcc 600
 c 601

<210> 8
 <211> 199
 <212> PRT
 <213> HOMO SAPIEN

<400> 8
 Thr Arg Val Pro Val Glu Ser Cys Gly Gln Tyr Gln Ser Cys Gly Glu
 1 5 10 15
 Cys Leu Gly Ser Gly Asp Pro His Cys Gly Trp Cys Val Leu His Asn
 20 25 30
 Thr Cys Thr Arg Lys Glu Arg Cys Glu Arg Ser Lys Glu Pro Arg Arg
 35 40 45
 Phe Ala Ser Glu Met Lys Gln Cys Val Arg Leu Thr Val His Pro Asn
 50 55 60
 Asn Ile Ser Val Ser Gln Tyr Asn Ala Leu Leu Val Leu Glu Thr Tyr
 65 70 75 80
 Asn Val Pro Glu Leu Ser Ala Gly Val Asn Cys Thr Phe Glu Asp Leu
 85 90 95
 Ser Glu Met Asp Gly Leu Val Val Gly Asn Gln Ile Gln Cys Tyr Ser
 100 105 110
 Pro Ala Ala Lys Glu Val Pro Arg Ile Ile Thr Glu Asn Gly Asp His
 115 120 125
 His Val Val Gln Leu Gln Leu Lys Ser Lys Glu Thr Gly Met Thr Phe
 130 135 140

Ala Ser Thr Ser Phe Val Phe Tyr Asn Cys Ser Val His Asn Ser Cys
 145 150 155 160
 Leu Ser Cys Val Glu Ser Pro Tyr Arg Cys His Trp Cys Lys Tyr Arg
 165 170 175
 His Val Cys Thr Asp Pro Lys Thr Cys Ser Phe Gln Glu Gly Arg Val
 180 185 190
 Lys Leu Pro Glu Val Gly Pro
 195

<210> 9
 <211> 6408
 <212> DNA
 <213> HOMO SAPIEN

<400> 9
 atgcctgtct tgggcccagc tcttctccag gctctctggg cggggtgggt cctcaccctc 60
 cagccctctc caccaaactgc attcaactccc aatggcacgt atctgcagca cctggcaagg 120
 gaccaccact caggcaccct ctaccctgggg gctaccaact tctctgtcca gctgagccct 180
 gggctagcag tggaggccac agtgctccac ggccctgtgc tagacagcag ggaactgcct 240
 ccactctgga tgcctgatga gtgccccccag gccacagccta ccaacaacccc gaatcagctg 300
 ctctctggga gccacggggc cctgggtgta tgcgggagcg tgcaccagggg tctctgtgaa 360
 cagcgggcgc tggggcagct cgagcagctg ctgctgcggc cagagcggccc tggggacaca 420
 caatagttgg tgcaccaatg tctctgcgtc agcacgglgg ggctggtagc ccaggggctt 480
 gcagggggag cctctctgtt tgggggggga ggatacacca gcagggggtgt ggggggtggc 540
 attccaccca tcacaacccc ggccctgtgg cgcccgacc ccgaagctgc ctctctctat 600
 gaggagacag ccaagctggc agtgggcgcg ctctccgagt accagcaca ctctctgagt 660
 gctcttgac ctggggccag cgtctacttc ctgttctgc ggcgggacct gcaggctcag 720
 tctagagctt ttctgctta tgtatctcga gtgtgtctcc ggagaccagca ctactactcc 780
 tatgtggagt tgcctctgac ctgcgaaggt ggcgcctaag ggctgatcca ggctcagct 840
 gtgcacagct ccagggaagt ggcccatggg gaggtgctct ttgcagcttt ctctctggct 900
 gcacccccca ctgtgggcgg gcccccactg cgccctgctg ggcctcttgg agcctctgct 960
 ctctgtgct tccccctgga tgaggtggac cggcttgcta atcgcacgcg agatgcctgc 1020
 taacaccagg aggtctgtgt tgaggaatgg accgaggtgg cctacatcga gtatgatgtc 1080
 aattctgagt ttgcacagct gccagtggac accctggatg cttatccctg tggctcagac 1140
 cacacgcccc agccccatgg ccagcgggtc ccgctggaag ccacaccaat tctggagttg 1200
 ccaggatttc agctaacagc tgtggcagtc accatggaag atggcacacac catcgctttc 1260
 ctgggtgata gtcaaggcca gctgcacagg gtctacttgg gccccaggag cgatggccac 1320
 ccatactcca cacagagagg ctcagcaggg tctgcagtga cagagacact cacctttgat 1380
 gggagctttg agcaccttga ttgctatgac cagagcacac ttctgaaggt tctgtggct 1440
 tctgtgctgt ccagccttga ctgtgcattc tgccctgtct acagggacccc atactctggg 1500
 tggctcgtgc tctctggcag gtgcagctgc cgttctgagt gctcgagggg ccagggccca 1560
 ggcagctggc tctggagctt ccagcctgag ctgggctgtc tgcaagtggc agccatgagt 1620
 cctgccaaca tcagccgaga ggagacgagg gaggttttcc tatcagtgcc agactctgcca 1680
 ccctgtggc caggggagtc atattctctg cactttgggg aacatcagag tctctccctg 1740
 ctgactgtgt ctggtgtgat gtgccccctc ccagacccta gtgagggccc agtgcgtccg 1800
 agaggagccg actacgtatc cgtgagcgtg gagctcagat ttggcgctgt tgtgatcgcc 1860
 aaaaactccc tctctttcta tgactgtgtg gcggtcactg aactccgccc atctgcgcag 1920
 tgcaggcgtg gtgtgagcag ccgctggggg tctaactggg gtgtctggca gcactctgtc 1980
 acccaacagg cctcgtgtga tgcctggccc atggttgcaa gccatcagag cccgcttgtc 2040
 tccccagac ctctctgaag aggttgacc agccccctcc caccacagc ccccaagacc 2100
 ctggccaccc ctgctctcga cacccttccc gtggagcctg gggctccctc cacagcaca 2160
 gctctggaca tctcaactgc ggctagtctc tccctgtcca gccctggggg gccatgggca 2220
 ggttctggct ccatacttc ccttggtctc acagggtcgc ctctccatga ggagccctcc 2280
 cctccagcag cccaaaatgg acctggaacc gctgtccctg ccccactga cttcagacc 2340
 tcagccacac ctgaggacct ctgtgcctcc ccgctgtcac cgtcagaggt agcagcagt 2400

ccccctgcag	accctggccc	cgaggetctt	catcccacag	tgccccctgga	cttggccccct	2460
gccactgttcc	ctgccaaccac	tttcccagg	gccatgggct	ccgtgaagcc	cgccctggag	2520
tggctcaaga	gagaaggcgg	cgagctgccc	gaggcggagc	agtggacggg	gggtgacgca	2580
cccgctctct	ccacttccac	ccctctctca	ggatgatggag	actcagcaga	gcttggagggc	2640
ctctccggccc	ccctcatctct	cccgctccagc	ctcgactacc	agtatgacac	ccccgggctc	2700
tgggagctgg	aagaggcgac	cttgggggca	agctctctgcc	ccgtgtgtgga	gagcgttcag	2760
ggctccacgt	tgatgccgtt	ccatgtggag	cgggaaatcc	ggctgctagg	caggaaacctg	2820
cacctttttcc	aggatggccc	aggagacaat	gagtggtgta	tggagctgga	gggctccgag	2880
gtggcgggtt	aggcccggtt	cgagtgtgag	ccacctccag	ataccagtg	ccatgtccacc	2940
tggccagcagc	accagctcag	ctatgaggct	ctgcagccgg	agctccgtgt	ggggctgttt	3000
ctgcgtatgg	cggtccgtct	gcgtgtggac	agtgtcagg	ggctgcattgt	ggatctgtat	3060
gactgttccg	tgggacattc	agactgcagc	cgctgccaaa	ctgccatgcc	ccagttatggc	3120
tgtgtgtgtt	gtgaggggga	gcgtccacgt	tgtgtgaaac	gggagggctg	tgttgaggctg	3180
gaggctgttg	ccaccacgtg	cccagcgccc	ctcatccact	cggtggagcc	actgactggg	3240
ctgtagacg	gaggcacccg	tgtaccatc	aggggctcca	acctggggcca	gcatgtgcag	3300
gatgtgtctg	gcattgtcac	gggtggctgga	gtgcccttgt	ctgtggatgc	ccaggagtatc	3360
gagggtctcca	cgagcctcgt	gtgcataccc	ggggccagtg	gggaggaggt	ggcccgggcc	3420
acagcgggtg	aggtgtccgg	aagaggagct	gggtgtccag	aacacgactt	tgctccacag	3480
gatccgaagg	tccattccat	cttcccggcc	cgccggccca	gagctggggg	accctgctctc	3540
accctgaattg	gctccaagct	ccctgactgg	cggtgtggag	acatccagat	gggtgttggga	3600
gaccagacgt	gtcacttgtc	gcgggagcag	cagtcagaac	aatcgtcggt	tgagacagc	3660
ccacggcccca	cgctgtccac	gtctcctgtg	gcgtgtgtgt	ttggggccac	ggagcggagg	3720
cttcaacgct	gacagttcaa	gtataacctg	gaccccaaca	tcacctctgc	tggtcccacc	3780
aagagcttccc	tcagttggag	acgttagata	tgcgtccgtg	gcagaaatct	ggagctggta	3840
cagacgcca	gaattccggg	gaccgtggtc	tcgagaatgc	ccaggcccaa	ccaggggctt	3900
ggacggaggc	tgcgtgttgt	cccggagacg	gcattgtccc	ttggaccctc	ctgcagtagc	3960
cagcaatttg	aggagccgtg	ccatgtcaac	tcctcccagc	tcataccgtg	cgcacacct	4020
gccctcccag	gcctgctctg	ggaccctctg	gtccgggtgg	aatttatctt	tgacaacctg	4080
gtctttgact	tgcacaactc	gaaccccaca	cttcttctct	atgaggccga	ccccaccctg	4140
cagccaactca	accctgagga	ccccaccatg	ccattccggc	ccattccctg	gagttgtgtc	4200
ctcgtgtgag	gggagaacct	ggacctgtga	atgtccaagg	aggagtggtg	ggctatgata	4260
ggggatggcc	ccctgtgtgt	gaagacgtca	acgcggccac	acctgtactg	cgagccccc	4320
gtggagcagc	ccctgccacc	gcaccatgcc	ctccagaggg	caactgactc	tttgcctgag	4380
ttcacctgtc	agatggggaa	cttgcgtctc	tccttgggtc	acgtgcagta	tgacggcgag	4440
agccctgggt	cttttctctg	ggcagccagc	gtgggcttgg	gggtggggac	ctctctctctg	4500
ctctgggtgt	tcatcatcat	tgctctcatg	tacaggagga	agagcaagca	ggccctagg	4560
gactataaag	aggttcagat	ccagctggag	aatctggaga	gcagtgtgct	ggaccctctg	4620
aagaaggaat	tcacagacct	catgactgag	atgaccgatc	tcaccagta	ctctctgggc	4680
agcgcatccc	ccctctctga	ctacaaggtg	tatgaggaga	ggatcttctt	ccctgggacc	4740
cgcgagctgc	cccttcacgc	ggacctgggt	gtgcctgaga	gcagacggcc	ccatgtagag	4800
caaggctctg	ggagactctc	taacctgtct	aacagcaagc	tcttctccac	caagtctcat	4860
ccacagctgc	agaccacagc	caccttttca	gtcctgggac	gtgcctacgt	ggcatctctg	4920
ctacccctgt	cactgcattg	gaagcttgag	tatttcaact	acatctctcg	cactctgctc	4980
agtgcagctg	tgtcccagta	tgtggccaa	aaecccaagc	tgatgctgcg	caggagacag	5040
acgtgtgttg	agaagctctc	caccaactgg	atgtccatct	gtctgtatac	ctctgtgagg	5100
gactccgtag	gggagctctc	gtacatgctc	tttcgaggga	ttaagcacc	agtggataag	5160
ggggcagctg	acagttgtgac	aggcaaggcc	aaatcacact	tgaaacgaca	cgctgctctc	5220
agagagagat	tggagtaccg	ctcccgtacc	ttgaatgcac	tattggctgt	ggggctctgg	5280
cgaggagag	cccaggcgct	gcccgtgaag	gtcttagact	gtgacacatc	ctcccaggca	5340
aagagagaag	tgtctgaaca	gctttataaa	ggagtgcctc	tcaccagcgc	gccagacctc	5400
gcaccccttg	atgtttagttg	gcggtctctgg	gtggccgggc	acctcatctt	tcttgacgag	5460
cagtctactt	ctcaggttcca	gggtctgtgg	aggcgcctga	acacactgca	gcattacaag	5520
gtcccagatg	gagcaactgt	ggccctctgt	cctgcctca	ccaagcatgt	gtccggggaa	5580
aaccagatct	atgtccctgt	agagcgacc	ccaatgctgg	aggatgtaga	tgaggggggc	5640
atccggccct	ggcacctggt	gaagccaagt	gatgagccgg	agccgcccag	gcctcggagg	5700

ggcagccttc	ggggcgggga	gcgtgagcgc	gccaaaggcca	tccttgagat	ctacctgacc	5760
cgccctgctg	ccatgaagg	caccctgcag	aagttcgtag	atgacctgtt	ccaggtgatt	5820
ctcagacaca	gccgccccgt	gccgctcgct	gtgaagtact	tctttgacct	gctggatgag	5880
caggccacgc	agcatggcat	ctccgaccag	gacacatccc	acatctggaa	gaccaacagc	5940
ttgctctcga	gggtctggat	caatataata	aaaaaccgcg	agtttggtgt	cgagctgcaa	6000
acatctgata	acatggatgc	ggtgctcctt	gtcattgcac	agaccttcat	ggagcgcctgc	6060
accttggcgc	accacaagct	gggcggggac	tccccgatca	acaaactctc	gtatgcaagg	6120
gacattcccc	ggtaacaagg	gatggtgga	aggtactatg	cagacatcag	acagactgtc	6180
ccagccagcg	accaagagat	gaactctgtc	ctggctgaac	tgctctggaa	ctactccgga	6240
gacctcgggg	cgcgagtgcc	cctgcatgaa	ctctacaagt	acatcaacaa	gtactatgac	6300
cagatcatca	ctgcctcgtg	ggaggatggc	acggcccaga	agatgcagct	gggtatcgg	6360
ctccagcaga	ttgcagctgc	tgtggaaaac	aaggtcacag	atctatag		6408

<210> 10

<211> 2135

<212> PRT

<213> HOMO SAPIEN

<400> 10

Met	Pro	Ala	Leu	Gly	Pro	Ala	Leu	Leu	Gln	Ala	Leu	Trp	Ala	Gly	Trp
1				5					10					15	
Val	Leu	Thr	Leu	Gln	Pro	Leu	Pro	Pro	Thr	Ala	Phe	Thr	Pro	Asn	Gly
			20					25					30		
Thr	Tyr	Leu	Gln	His	Leu	Ala	Arg	Asp	Pro	Thr	Ser	Gly	Thr	Leu	Tyr
			35				40					45			
Leu	Gly	Ala	Thr	Asn	Phe	Leu	Phe	Gln	Leu	Ser	Pro	Gly	Leu	Gln	Leu
			50			55					60				
Glu	Ala	Thr	Val	Ser	Thr	Gly	Pro	Val	Leu	Asp	Ser	Arg	Asp	Cys	Leu
			65			70				75				80	
Pro	Pro	Val	Met		Pro	Asp	Glu	Cys	Pro	Gln	Ala	Gln	Pro	Thr	Asn
			85						90					95	
Pro	Asn	Gln	Leu	Leu	Leu	Val	Ser	Pro	Gly	Ala	Leu	Val	Val	Cys	Gly
			100					105					110		
Ser	Val	His	Gln	Gly	Val	Cys	Glu	Gln	Arg	Arg	Leu	Gly	Gln	Leu	Glu
			115				120					125			
Gln	Leu	Leu	Leu	Arg	Pro	Glu	Arg	Pro	Gly	Asp	Thr	Gln	Tyr	Val	Ala
			130			135					140				
Ala	Asn	Asp	Pro	Ala	Val	Ser	Thr	Val	Gly	Leu	Val	Ala	Gln	Gly	Leu
			145			150				155				160	
Ala	Gly	Glu	Pro	Leu	Leu	Phe	Val	Gly	Arg	Gly	Tyr	Thr	Ser	Arg	Gly
			165					170					175		
Val	Gly	Gly	Gly	Ile	Pro	Pro	Ile	Thr	Thr	Arg	Ala	Leu	Trp	Pro	Pro
			180					185					190		
Asp	Pro	Gln	Ala	Ala	Phe	Ser	Tyr	Glu	Glu	Thr	Ala	Lys	Leu	Ala	Val
			195				200				205				
Gly	Arg	Leu	Ser	Glu	Tyr	Ser	His	His	Phe	Val	Ser	Ala	Phe	Ala	Arg
			210			215					220				
Gly	Ala	Ser	Ala	Tyr	Phe	Leu	Phe	Leu	Arg	Arg	Asp	Leu	Gln	Ala	Gln
			225			230				235				240	
Ser	Arg	Ala	Phe	Arg	Ala	Tyr	Val	Ser	Arg	Val	Cys	Leu	Arg	Asp	Gln
			245					250					255		
His	Tyr	Tyr	Ser	Tyr	Val	Glu	Leu	Pro	Leu	Ala	Cys	Glu	Gly	Gly	Arg
			260				265						270		

Tyr Gly Leu Ile Gln Ala Ala Ala Val Ala Thr Ser Arg Glu Val Ala
 275 280
 His Gly Glu Val Leu Phe Ala Ala Phe Ser Ser Ala Ala Pro Pro Thr
 290 295
 Val Gly Arg Pro Pro Ser Ala Ala Ala Gly Ala Ser Gly Ala Ser Ala
 305 310 315 320
 Leu Cys Ala Phe Pro Leu Asp Glu Val Asp Arg Leu Ala Asn Arg Thr
 325 330 335
 Arg Asp Ala Cys Tyr Thr Arg Glu Gly Arg Ala Glu Asp Gly Thr Glu
 340 345 350
 Val Ala Tyr Ile Glu Tyr Asp Val Asn Ser Asp Cys Ala Gln Leu Pro
 355 360 365
 Val Asp Thr Leu Asp Ala Tyr Pro Cys Gly Ser Asp His Thr Pro Ser
 370 375 380
 Pro Met Ala Ser Arg Val Pro Leu Glu Ala Thr Pro Ile Leu Glu Trp
 385 390 395 400
 Pro Gly Ile Gln Leu Thr Ala Val Ala Val Thr Met Glu Asp Gly His
 405 410 415
 Thr Ile Ala Phe Leu Gly Asp Ser Gln Gly Gln Leu His Arg Val Tyr
 420 425 430
 Leu Gly Pro Gly Ser Asp Gly His Pro Tyr Ser Thr Gln Ser Ile Gln
 435 440 445
 Gln Gly Ser Ala Val Ser Arg Asp Leu Thr Phe Asp Gly Thr Phe Glu
 450 455 460
 His Leu Tyr Val Met Thr Gln Ser Thr Leu Leu Lys Val Pro Val Ala
 465 470 475 480
 Ser Cys Ala Gln His Leu Asp Cys Ala Ser Cys Leu Ala His Arg Asp
 485 490 495
 Pro Tyr Cys Gly Trp Cys Val Leu Leu Gly Arg Cys Ser Arg Arg Ser
 500 505 510
 Glu Cys Ser Arg Gly Gln Gly Pro Glu Gln Trp Leu Trp Ser Phe Gln
 515 520 525
 Pro Glu Leu Gly Cys Leu Gln Val Ala Ala Met Ser Pro Ala Asn Ile
 530 535 540
 Ser Arg Glu Glu Thr Arg Glu Val Phe Leu Ser Val Pro Asp Leu Pro
 545 550 555 560
 Pro Leu Trp Pro Gly Glu Ser Tyr Ser Cys His Phe Gly Glu His Gln
 565 570 575
 Ser Pro Ala Leu Leu Thr Gly Ser Gly Val Met Cys Pro Ser Pro Asp
 580 585 590
 Pro Ser Glu Ala Pro Val Leu Pro Arg Gly Ala Asp Tyr Val Ser Val
 595 600 605
 Ser Val Glu Leu Arg Phe Gly Ala Val Val Ile Ala Lys Thr Ser Leu
 610 615 620
 Ser Phe Tyr Asp Cys Val Ala Val Thr Glu Leu Arg Pro Ser Ala Gln
 625 630 635 640
 Cys Gln Ala Cys Val Ser Ser Arg Trp Gly Cys Asn Trp Cys Val Trp
 645 650 655
 Gln His Leu Cys Thr His Lys Ala Ser Cys Asp Ala Gly Pro Met Val
 660 665 670
 Ala Ser His Gln Ser Pro Leu Val Ser Pro Asp Pro Pro Ala Arg Gly
 675 680 685
 Gly Pro Ser Pro Ser Pro Pro Thr Ala Pro Lys Ala Leu Ala Thr Pro
 690 695 700
 Ala Pro Asp Thr Leu Pro Val Glu Pro Gly Ala Pro Ser Thr Ala Thr

705				710				715				720
Ala Ser Asp Ile	Ser	Pro Gly Ala Ser	Pro	Ser Leu Leu Ser	Pro							Trp
Gly Pro Trp Ala	725	Gly Ser Gly Ser	730	Ser Ser Pro Gly Ser	735							Gly
	740		745		750							
Ser Pro Leu His	Glu	Glu Pro Ser Pro	Pro	Ser Pro Gln Asn Gly	755							Pro
Gly Thr Ala Val	Pro	Ala Pro Thr Asp	Phe	Arg Pro Ser Ala Thr	760							Pro
	770		775		780							
Glu Asp Leu Leu	Ala	Ser Pro Leu Ser	Pro	Ser Glu Val Ala Ala	785							Val
Pro Pro Ala Asp	790	Pro Gly Pro Glu Ala	Leu	His Pro Thr Val Pro	800							Leu
	805		810		815							
Asp Leu Pro Pro	Ala	Thr Val Pro Ala	Thr	Phe Pro Gly Ala Met	820							
	825		830		835							
Gly Ser Val Lys	Pro	Ala Leu Asp Trp	Leu	Thr Arg Glu Gly Gly	840							Glu
	845		850		855							
Leu Pro Glu Ala	Asp	Glu Trp Thr Gly	Gly	Asp Ala Pro Ala Phe	860							Ser
Thr Ser Thr Leu	Leu	Ser Gly Asp Gly	Asp	Ser Ala Glu Leu Glu	865							Gly
	870		875		880							
Pro Pro Ala Pro	Leu	Ile Leu Pro Ser	Ser	Leu Asp Tyr Gln Tyr	885							Asp
	890		895		900							
Thr Pro Gly Leu	Trp	Glu Leu Glu Glu	Ala	Thr Leu Gly Ala Ser	905							Ser
	910		915		920							
Cys Pro Cys Val	Glu	Ser Val Gln Gly	Ser	Thr Leu Met Pro	925							Val
	930		935		940							
Val Glu Arg Glu	Ile	Arg Leu Leu Gly	Arg	Asn Leu His Leu	945							Phe
	950		955		960							
Asp Gly Pro Gly	Asp	Asn Glu Cys Val	Met	Glu Leu Glu Gly Leu	965							Glu
Val Val Val Glu	Ala	Arg Val Glu Cys	Glu	Pro Pro Asp Thr	970							Gln
	975		980		985							
Cys His Val Thr	Cys	Gln Gln His Gln	Leu	Ser Tyr Glu Ala Leu	990							Gln
	995		1000		1005							
Pro Glu Leu Arg	Val	Gly Leu Phe Leu	Arg	Arg Ala Gly Arg Leu	1010							Arg
	1015		1020		1025							
Val Asp Ser Ala	Glu	Gly Leu His Val	Val	Leu Tyr Asp Cys	1030							Val
	1035		1040		1045							
Gly His Gly Asp	Cys	Ser Arg Cys Gln	Thr	Ala Met Pro Gln Tyr	1050							Gly
	1055		1060		1065							
Cys Val Trp Cys	Glu	Gly Glu Arg Pro	Arg	Cys Val Thr Arg Glu	1070							Ala
	1075		1080		1085							
Cys Gly Glu Ala	Glu	Ala Val Ala Thr	Gln	Cys Pro Ala Pro Leu	1090							Ile
	1095		1100		1105							
His Ser Val Glu	Pro	Leu Thr Gly Pro	Val	Asp Gly Gly Thr Arg	1110							Val
	1115		1120		1125							
Thr Ile Arg Gly	Ser	Asn Leu Gly Gln	His	Val Gln Asp Val Leu	1130							Gly
	1135		1140		1145							
Met Val Thr Val	Ala	Gly Val Pro Cys	Ala	Val Asp Ala Gln Glu	1150							Tyr
	1155		1160		1165							
Glu Val Ser Ser	Ser	Leu Val Cys Ile	Thr	Gly Ala Ser Gly Glu	1170							Glu
	1175		1180		1185							
Val Ala Gly Ala	Thr	Ala Val Glu Val	Pro	Gly Arg Gly Arg Gly	1190							Val
	1195		1200		1205							

Ser Glu His Asp Phe Ala Tyr Gln Asp Pro Lys Val His Ser Ile Phe
 1155 1160 1165
 Pro Ala Arg Gly Pro Arg Ala Gly Gly Thr Arg Leu Thr Leu Asn Gly
 1170 1175 1180
 Ser Lys Leu Leu Thr Gly Arg Leu Glu Asp Ile Arg Val Val Val Gly
 1185 1190 1195 1200
 Asp Gln Pro Cys His Leu Leu Pro Glu Gln Gln Ser Glu Gln Leu Arg
 1205 1210 1215
 Cys Glu Thr Ser Pro Arg Pro Thr Pro Ala Thr Leu Pro Val Ala Val
 1220 1225 1230
 Trp Phe Gly Ala Thr Glu Arg Arg Leu Gln Arg Gly Gln Phe Lys Tyr
 1235 1240 1245
 Thr Leu Asp Pro Asn Ile Thr Ser Ala Gly Pro Thr Lys Ser Phe Leu
 1250 1255 1260
 Ser Gly Gly Arg Glu Ile Cys Val Arg Gly Gln Asn Leu Asp Val Val
 1265 1270 1275 1280
 Gln Thr Pro Arg Ile Arg Val Thr Val Val Ser Arg Met Leu Gln Pro
 1285 1290 1295
 Ser Gln Gly Leu Gly Arg Arg Arg Arg Val Val Pro Glu Thr Ala Cys
 1300 1305 1310
 Ser Leu Gly Pro Ser Cys Ser Ser Gln Gln Phe Glu Glu Pro Cys His
 1315 1320 1325
 Val Asn Ser Ser Gln Leu Ile Thr Cys Arg Thr Pro Ala Leu Pro Gly
 1330 1335 1340
 Leu Pro Glu Asp Pro Trp Val Arg Val Glu Phe Ile Leu Asp Asn Leu
 1345 1350 1355 1360
 Val Phe Asp Phe Ala Thr Leu Asn Pro Thr Pro Phe Ser Tyr Glu Ala
 1365 1370 1375
 Asp Pro Thr Leu Gln Pro Leu Asn Pro Glu Asp Pro Thr Met Pro Phe
 1380 1385 1390
 Arg His Lys Pro Gly Ser Val Phe Ser Val Glu Gly Glu Asn Leu Asp
 1395 1400 1405
 Leu Ala Met Ser Lys Glu Glu Val Val Ala Met Ile Gly Asp Gly Pro
 1410 1415 1420
 Cys Val Val Lys Thr Leu Thr Arg His His Leu Tyr Cys Glu Pro Pro
 1425 1430 1435 1440
 Val Glu Gln Pro Leu Pro Arg His His Ala Leu Arg Glu Ala Pro Asp
 1445 1450 1455
 Ser Leu Pro Glu Phe Thr Val Gln Met Gly Asn Leu Arg Phe Ser Leu
 1460 1465 1470
 Gly His Val Gln Tyr Asp Gly Glu Ser Pro Gly Ala Phe Pro Val Ala
 1475 1480 1485
 Ala Gln Val Gly Leu Gly Val Gly Thr Ser Leu Leu Ala Leu Gly Val
 1490 1495 1500
 Ile Ile Ile Val Leu Met Tyr Arg Arg Lys Ser Lys Gln Ala Leu Arg
 1505 1510 1515 1520
 Asp Tyr Lys Lys Val Gln Ile Gln Leu Glu Asn Leu Glu Ser Ser Val
 1525 1530 1535
 Arg Asp Arg Cys Lys Lys Glu Phe Thr Asp Leu Met Thr Glu Met Thr
 1540 1545 1550
 Asp Leu Thr Ser Asp Leu Leu Gly Ser Gly Ile Pro Phe Leu Asp Tyr
 1555 1560 1565
 Lys Val Tyr Ala Glu Arg Ile Phe Phe Pro Gly His Arg Glu Ser Pro
 1570 1575 1580
 Leu His Arg Asp Leu Gly Val Pro Glu Ser Arg Arg Pro Thr Val Glu

1585					1590					1595					1600
Gln	Gly	Leu	Gly	Gln	Leu	Ser	Asn	Leu	Leu	Asn	Ser	Lys	Leu	Phe	Leu
				1605						1610					1615
Thr	Lys	Phe	Ile	His	Thr	Leu	Glu	Ser	Gln	Arg	Thr	Phe	Ser	Ala	Arg
				1620						1625					1630
Asp	Arg	Ala	Tyr	Val	Ala	Ser	Leu	Leu	Thr	Val	Ala	Leu	His	Gly	Lys
				1635						1640					1645
Leu	Glu	Tyr	Phe	Thr	Asp	Ile	Leu	Arg	Thr	Leu	Leu	Ser	Asp	Leu	Val
				1650						1655					1660
Ala	Gln	Tyr	Val	Ala	Lys	Asn	Pro	Lys	Leu	Met	Leu	Arg	Arg	Thr	Glu
				1665						1670					1675
Thr	Val	Val	Glu	Lys	Leu	Leu	Thr	Asn	Trp	Met	Ser	Ile	Cys	Leu	Tyr
				1685						1690					1695
Thr	Phe	Val	Arg	Asp	Ser	Val	Gly	Glu	Pro	Leu	Tyr	Met	Leu	Phe	Arg
				1700						1705					1710
Gly	Ile	Lys	His	Gln	Val	Asp	Lys	Gly	Pro	Val	Asp	Ser	Val	Thr	Gly
				1715						1720					1725
Lys	Ala	Lys	Tyr	Thr	Leu	Asn	Asp	Asn	Arg	Leu	Leu	Arg	Glu	Asp	Val
				1730						1735					1740
Glu	Tyr	Arg	Pro	Leu	Thr	Leu	Asn	Ala	Leu	Leu	Ala	Val	Gly	Pro	Gly
				1745						1750					1755
Ala	Gly	Glu	Ala	Gln	Gly	Val	Pro	Val	Lys	Val	Leu	Asp	Cys	Asp	Thr
				1765						1770					1775
Ile	Ser	Gln	Ala	Lys	Glu	Lys	Met	Leu	Asp	Gln	Leu	Tyr	Lys	Gly	Val
				1780						1785					1790
Pro	Leu	Thr	Gln	Arg	Pro	Asp	Pro	Arg	Thr	Leu	Asp	Val	Glu	Trp	Arg
				1795						1800					1805
Ser	Gly	Val	Ala	Gly	His	Leu	Ile	Leu	Ser	Asp	Glu	Asp	Val	Thr	Ser
				1810						1815					1820
Glu	Val	Gln	Gly	Leu	Trp	Arg	Arg	Leu	Asn	Thr	Leu	Gln	His	Tyr	Lys
				1825						1830					1835
Val	Pro	Asp	Gly	Ala	Thr	Val	Ala	Leu	Val	Pro	Cys	Leu	Thr	Lys	His
				1845						1850					1855
Val	Leu	Arg	Glu	Asn	Gln	Asp	Tyr	Val	Pro	Gly	Glu	Arg	Thr	Pro	Met
				1860						1865					1870
Leu	Glu	Asp	Val	Asp	Glu	Gly	Gly	Ile	Arg	Pro	Trp	His	Leu	Val	Lys
				1875						1880					1885
Pro	Ser	Asp	Glu	Pro	Glu	Pro	Pro	Arg	Pro	Arg	Arg	Gly	Ser	Leu	Arg
				1890						1895					1900
Gly	Gly	Glu	Arg	Glu	Arg	Ala	Lys	Ala	Ile	Pro	Glu	Ile	Tyr	Leu	Thr
				1905						1910					1915
Arg	Leu	Leu	Ser	Met	Lys	Gly	Thr	Leu	Gln	Lys	Phe	Val	Asp	Asp	Leu
				1925						1930					1935
Phe	Gln	Val	Ile	Leu	Ser	Thr	Ser	Arg	Pro	Val	Pro	Leu	Ala	Val	Lys
				1940						1945					1950
Tyr	Phe	Phe	Asp	Leu	Leu	Asp	Glu	Gln	Ala	Gln	Gln	His	Gly	Ile	Ser
				1955						1960					1965
Asp	Gln	Asp	Thr	Ile	His	Ile	Trp	Lys	Thr	Asn	Ser	Leu	Pro	Leu	Arg
				1970						1975					1980
Phe	Trp	Ile	Asn	Ile	Ile	Lys	Asn	Pro	Gln	Phe	Val	Phe	Asp	Val	Gln
				1985						1990					1995
Thr	Ser	Asp	Asn	Met	Asp	Ala	Val	Leu	Leu	Val	Ile	Ala	Gln	Thr	Phe
				2005						2010					2015
Met	Asp	Ala	Cys	Thr	Leu	Ala	Asp	His	Lys	Leu	Gly	Arg	Asp	Ser	Pro
				2020						2025					2030

Ile Asn Lys Leu Leu Tyr Ala Arg Asp Ile Pro Arg Tyr Lys Arg Met
 2035 2040 2045
 Val Glu Arg Tyr Tyr Ala Asp Ile Arg Gln Thr Val Pro Ala Ser Asp
 2050 2055 2060
 Gln Glu Met Asn Ser Val Leu Ala Glu Leu Ser Trp Asn Tyr Ser Gly
 2065 2070 2075 2080
 Asp Leu Gly Ala Arg Val Ala Leu His Glu Leu Tyr Lys Tyr Ile Asn
 2085 2090 2095
 Lys Tyr Tyr Asp Gln Ile Ile Thr Ala Leu Glu Glu Asp Gly Thr Ala
 2100 2105 2110
 Gln Lys Met Gln Leu Gly Tyr Arg Leu Gln Gln Ile Ala Ala Ala Val
 2115 2120 2125
 Glu Asn Lys Val Thr Asp Leu
 2130 2135

<210> 11

<211> 2190

<212> DNA

<213> HOMO SAPIEN

<400> 11

atgctctgctc	tgggcccagc	tcttctccag	gctctctggg	ccgggtgggt	cctcacccctc	60
cagcccccctc	caccactctc	atctactccc	aatggcacgt	atctgcagca	cctggcgaagg	120
cagcccaactc	caggcacctc	ctactctggg	gctaccaact	tctgtttcca	gctgagccct	180
gggctgagc	tggaggccac	agtgtccacc	ggccctgtgc	tagacagcag	ggactgcctg	240
ccactctga	tgcctgatga	gtgccccag	gccagcccta	ccaacaacc	gaatcagctg	300
ctctcgtgtg	gccccagggc	cctgtgtgta	tgcgggagcg	tgcaccaggg	ggctgtgtga	360
cagcggcgcc	tggggcagct	cgagcagctg	ctgctgcggc	cagagcggcc	tggggacaca	420
caatatgtgg	ctgccaatga	tctctcggtc	agcacgggtg	ggctggtagc	ccagggcctg	480
gcaggggagc	ccctcctgtt	tgtggggcga	ggatacacca	gcagggggtg	ggggggtggc	540
attccaccaca	tcacaacccc	ggccctgtgg	ccgcccgacc	cccaagctgc	cttctcctat	600
gagggagacg	ccaagctggc	agtgggcgcg	ctctccgagt	acagccacca	cttcgtgagt	660
gcctttgcac	gtggggccag	cgctacttct	ctgttctctc	ggcgggacct	gcaggctcag	720
tctagagctt	tctgtgccta	tgtatctcga	gtgtgtctcc	gggaccagca	ctactactcc	780
tatgtggagt	tgcctctggc	ctgcgaaggt	ggccgctacg	ggctgatcca	ggctgcagct	840
gtggccacgt	ccaggggagt	ggcgcatggg	gaggtgctct	ttgcagcttt	ctctctggct	900
gcacccccca	ctgtgggcgc	gccccatcgc	gcggctgctg	gggcatctgg	agcctctgcc	960
ctctgtgcct	tccccctgga	tgaggtggac	cggttgctga	atcgcacgcg	agatgcctctg	1020
tacacccggg	agggtcgtgc	tgaggatggg	accgaggtgg	cctacatcga	gtatgatgtc	1080
aattctgact	gtgcacagct	gccagtgagc	accctggatg	cttatccctg	tgtctcagac	1140
cacacgcccc	gccccatggc	cagccgggtc	ccgctggaag	ccacaccaat	tctggagtgg	1200
ccagggatcc	agctaacacg	tgtggcagtc	accatggaag	atggacacac	catctcttct	1260
ctgggtgata	gtcaaggcca	gctgcacagc	gtctacttgg	gccacgggag	cgatggccac	1320
ccatactaca	cacagagcat	ccagcagggg	tctgcagtga	gcagagacct	cacctttgat	1380
gggacctttg	agcacctgta	tgtcatgacc	cagagcacac	ttctgaaggt	tcctgtggct	1440
tcctctgtctc	agcacctgga	ctgtgcacat	tgccttgctc	acagggaccc	atactgtggg	1500
tggtgcgtgc	tccttggcag	gtgcagtcgc	cgttctgagt	gctcaggggg	ccagggccca	1560
gagcagtgcc	tatggagctt	ccagcctgag	ctgggctgtc	tgcgaagtgc	agccatgagt	1620
cttgccaaca	tcagccgaga	ggagacgagg	gaggttttcc	tatcagtgcc	agacctgcca	1680
ccctctgtgc	caggggagtc	ataattctgc	cactttgggg	aacatcagag	atctgccctg	1740
ctgacttggt	ctggtgtgat	gtgccccctc	ccagacccta	gtgagggccc	agtgctgcgc	1800
agaggagccg	actacgtatc	ctgtgagcgt	gagctcagat	ttggcgctgt	tgtgatcgcc	1860
aaaaactccc	tctctttcta	tgaactgtgt	gcggctcact	aactccggcc	atctgcgcag	1920
tgccagcgct	gtgtgagcag	ccgctggggg	tgtaaactgg	gtgtctggca	gcacctgtgc	1980
acccacaaag	cctcgtgtga	tgtctgggcc	atggttgcaa	gccatcaggt	gatggagact	2040

cagcagagct	tgagggcct	ccgcgcctcc	tcctctccc	gtccagcctc	gactaccagt	2100
atgacacccc	cgggctctgg	gagctggaag	aggcgacctt	gggggcaagc	tctgtccctt	2160
gtgtggagag	cggtcagggc	tccacgttga				2190

<210> 12
 <211> 729
 <212> PRT
 <213> HOMO SAPIEN

<400> 12
 Met Pro Ala Leu Gly Pro Ala Leu Leu Gln Ala Leu Trp Ala Gly Trp
 1 5 10 15
 Val Leu Thr Leu Gln Pro Leu Pro Pro Thr Ala Phe Thr Pro Asn Gly
 20 25 30
 Thr Tyr Leu Gln His Leu Ala Arg Asp Pro Thr Ser Gly Thr Leu Tyr
 35 40 45
 Leu Gly Ala Thr Asn Phe Leu Phe Gln Leu Ser Pro Gly Leu Gln Leu
 50 55 60
 Glu Ala Thr Val Ser Thr Gly Pro Val Leu Asp Ser Arg Asp Cys Leu
 65 70 75 80
 Pro Pro Val Met Pro Asp Glu Cys Pro Gln Ala Gln Pro Thr Asn Asn
 85 90 95
 Pro Asn Gln Leu Leu Val Ser Pro Gly Ala Leu Val Val Cys Gly
 100 105 110
 Ser Val His Gln Gly Val Cys Glu Gln Arg Arg Leu Gly Gln Leu Glu
 115 120 125
 Gln Leu Leu Leu Arg Pro Glu Arg Pro Gly Asp Thr Gln Tyr Val Ala
 130 135 140
 Ala Asn Asp Pro Ala Val Ser Thr Val Gly Leu Val Ala Gln Gly Leu
 145 150 155 160
 Ala Gly Glu Pro Leu Leu Phe Val Gly Arg Gly Tyr Thr Ser Arg Gly
 165 170 175
 Val Gly Gly Gly Ile Pro Pro Ile Thr Thr Arg Ala Leu Trp Pro Pro
 180 185 190
 Asp Pro Gln Ala Ala Phe Ser Tyr Glu Glu Thr Ala Lys Leu Ala Val
 195 200 205
 Gly Arg Leu Ser Glu Tyr Ser His His Phe Val Ser Ala Phe Ala Arg
 210 215 220
 Gly Ala Ser Ala Tyr Phe Leu Phe Leu Arg Arg Asp Leu Gln Ala Gln
 225 230 235 240
 Ser Arg Ala Phe Arg Ala Tyr Val Ser Arg Val Cys Leu Arg Asp Gln
 245 250 255
 His Tyr Tyr Ser Tyr Val Glu Leu Pro Leu Ala Cys Glu Gly Gly Arg
 260 265 270
 Tyr Gly Leu Ile Gln Ala Ala Val Ala Thr Ser Arg Glu Val Ala
 275 280 285
 His Gly Glu Val Leu Phe Ala Ala Phe Ser Ser Ala Ala Pro Pro Thr
 290 295 300
 Val Gly Arg Pro Pro Ser Ala Ala Ala Gly Ala Ser Gly Ala Ser Ala
 305 310 315 320
 Leu Cys Ala Phe Pro Leu Asp Glu Val Asp Arg Leu Ala Asn Arg Thr
 325 330 335
 Arg Asp Ala Cys Tyr Thr Arg Glu Gly Arg Ala Glu Asp Gly Thr Glu
 340 345 350
 Val Ala Tyr Ile Glu Tyr Asp Val Asn Ser Asp Cys Ala Gln Leu Pro

[illegible]